

Is the five-flow conjecture almost false?

Jesper Lykke Jacobsen^{1,2} and Jesús Salas³

¹*Laboratoire de Physique Théorique, École Normale Supérieure
24 rue Lhomond, 75231 Paris, FRANCE
JESPER.JACOBSEN@ENS.FR*

²*Université Pierre et Marie Curie, 4 place Jussieu, 75252 Paris, FRANCE*

³*Instituto Gregorio Millán
and
Grupo de Modelización, Simulación Numérica y Matemática Industrial
Universidad Carlos III de Madrid
Avda. de la Universidad, 30
28911 Leganés, SPAIN
JSALAS@MATH.UC3M.ES*

October 19, 2010

Abstract

The number of nowhere-zero $\mathbb{Z}_Q$ flows on a graph G can be shown to be a polynomial in Q , defining the flow polynomial $\Phi_G(Q)$. According to Tutte's five-flow conjecture, $\Phi_G(5) > 0$ for any bridgeless G . A conjecture by Welsh that $\Phi_G(Q)$ has no real roots for $Q \in (4, \infty)$ was recently disproved by Haggard, Pearce and Royle. These authors conjectured the absence of roots for $Q \in [5, \infty)$. We study the real and complex roots of $\Phi_G(Q)$ for a family of non-planar cubic graphs known as generalised Petersen graphs $G(n, k)$. We show that the modified conjecture on real flow roots is also false, by exhibiting infinitely many real flow roots $Q > 5$ within the class $G(nk, k)$. In particular, we compute explicitly $G(119, 7)$ showing that it has real roots at $Q \approx 5.0000197675$ and $Q \approx 5.1653424423$. We moreover prove that the graph families $G(6n, 6)$ and $G(7n, 7)$ possess real flow roots that accumulate at $Q = 5$ as $n \rightarrow \infty$ (in the latter case from above and below); and that $Q_c(7) \approx 5.2352605291$ is an accumulation point of real zeros of the flow polynomials for $G(7n, 7)$ as $n \rightarrow \infty$.

Key Words: Nowhere-zero flows; flow polynomial; flow roots; Tutte's five-flow conjecture; Petersen graph; transfer matrix.

1 Introduction

The four-colour problem is probably the best known problem of graph theory, and has fascinated generations of mathematicians and laymen since it was first stated in a letter from Augustus de Morgan to Sir William Rowan Hamilton dated 23 October 1852. The suspense was resolved in late July 1976 when Appel and Haken [1] announced the proof of the four-colour theorem: Every planar graph admits a 4-vertex-colouring.

A quantitative version of the problem was proposed by Birkhoff [14] in 1912: Given an arbitrary graph G and a set of Q colours, how many proper vertex colourings does G admit? This defines the chromatic polynomial $\chi_G(Q)$. The four-colour theorem can then be stated: If G is planar, $\chi_G(4) > 0$.

Although initially defined for $Q \in \mathbb{N}$, it is easily proved [14, 58, 59] that $\chi_G(Q)$ is in fact a polynomial in Q , and can as such be evaluated for any $Q \in \mathbb{C}$. This suggests an algebraic or even analytic approach to the colouring problem. There exist many studies of the location in $\mathbb{C}$ of the roots of $\chi_G(Q)$, henceforth called chromatic roots. These studies concern either specific graphs, or all planar graphs, or other infinite families of graphs.

Birkhoff and Lewis [15] have made the following conjecture: If G is planar, $\chi_G(Q) > 0$ for $Q \in [4, \infty)$. Obviously, the statement of this conjecture is stronger than the four-colour theorem, but unfortunately has not yet been turned into a theorem. The corresponding result for $Q \in [5, \infty)$ has however been proved by the same authors [15] (see also [37, 50, 60]).

Beraha and Kahane [8] have exhibited an infinite family of planar graphs for which $Q = 4$ can be proved to be an accumulation point of *complex* chromatic roots. In that sense, the four-colour theorem is “almost false”. Improving on this, Royle [41] has proved, for a slightly different family, that $Q = 4$ is also an accumulation point of *real* chromatic roots (converging to $Q = 4$ from below). Finally, Sokal [48] has proved that for a specific family of planar graphs (generalised Θ -graphs) chromatic roots are dense in $\mathbb{C}$ (except perhaps in the disc $|Q - 1| < 1$).

A close cousin of the chromatic polynomial is the so-called *flow polynomial* $\Phi_G(Q)$. Let $G = (V, E)$ be an arbitrary (not necessarily planar) graph G with vertex set V and edge set E , and let Γ be an additive Abelian group. A Γ -flow on G is a map $\phi: E \mapsto \Gamma$ that attributes a variable $\phi(e)$ to each edge $e \in E$, subject to the conservation of these variables at each vertex, with respect to an arbitrary chosen orientation of E . An elementary example of a flow is the current ϕ in an electrical network, in which case the conservation constraint is known as Kirchhoff’s first law [34].

A *nowhere zero* Γ -flow is a Γ -flow ϕ such that $\phi(e) \neq 0$ for all $e \in E$ [33, 54, 63]. If Γ is a *finite* Abelian group of order Q , it can be shown that the number of nowhere zero Γ -flows depends only on Q (not on the *specific* structure of the group Γ), and it is in fact the restriction to $Q \in \mathbb{N}$ of a polynomial in Q called the flow polynomial $\Phi_G(Q)$ [52]. One can then again extend the definition to $Q \in \mathbb{C}$ and study the location of (real or complex) flow roots.

A *nowhere zero* Q -flow of G is a nowhere zero $\mathbb{Z}$ -flow ϕ such that $|\phi(e)| \leq |Q - 1|$ for all $e \in E$. Tutte [52] showed that G has a nowhere zero Q -flow if and only if it has a nowhere zero $\mathbb{Z}_Q$ -flow; but these two concepts are different! Tutte’s result immediately implies the following interesting (but far from obvious) property of nowhere-zero $\mathbb{Z}_Q$ -flows [53]:

Proposition 1.1 *If $\Phi_G(Q) > 0$ for some $Q \in \mathbb{N}$, then $\Phi_G(Q') > 0$ for all integers $Q' \geq Q$.*

When G is planar, one has [51] the duality relation $\chi_{G^*}(Q) = Q \Phi_G(Q)$, where G^* denotes the dual graph. The properties of $\Phi_G(Q)$ thus follow from those of $\chi_{G^*}(Q)$.

It is worth stressing that the Birkhoff-Lewis theorem [15] provides a uniform upper bound for the real zeros of the chromatic polynomial of *all* planar graphs, namely $Q = 5$. However, such an upper bound (if it actually exists at all!) is not known for the real zeros of the flow polynomial of arbitrary bridgeless graphs.¹ The existence of such uniform upper bound and its value, if it does exist, are long-standing open problems in Combinatorics.

Consider now arbitrary (not necessarily planar) bridgeless graphs G . Because there exist graphs not admitting a nowhere zero 4-flow,² the strongest possible results for integer and real flow roots are given respectively by the following two well-known conjectures:

Conjecture 1.2 (Tutte’s five-flow conjecture [33, 53, 54]) *For any bridgeless graph G , $\Phi_G(5) > 0$.*

Remark. This conjecture implies that $\Phi_G(Q) > 0$ for all integers $Q \geq 5$ by Proposition 1.1.

Conjecture 1.3 (Welsh [57]) *For any bridgeless graph G , $\Phi_G(Q) > 0$ for $Q \in (4, \infty)$.*

It should be noted that the Welsh conjecture parallels that of Birkhoff and Lewis for the chromatic polynomial. It does however not include the case $Q = 4$. Some results by Jackson on zero-free intervals for the flow polynomials of cubic graphs [24, 25] also suggest this close parallelism between $\chi_G(Q)$ for planar G and $\Phi_G(Q)$ for *arbitrary* G . (Note that both polynomials are evaluated at the *same* value of Q .)

A number of weaker results have been proved over the years, notably:

Theorem 1.4 (Seymour [46]) *For any bridgeless graph G , $\Phi_G(6) > 0$.*

Theorem 1.5 (Steinberg [49]) *For any bridgeless graph G that is embeddable in the projective plane, $\Phi_G(5) > 0$.*

An immediate Corollary of Seymour’s theorem (using Proposition 1.1) is that $\Phi_G(Q) > 0$ for all integers $Q \geq 6$. Thus $Q = 5$ is a uniform upper bound for *integer* flow roots. But the above results give no clue about the existence of a uniform upper bound for *real* flow roots.

A first step into proving (or disproving) Conjectures 1.2–1.3 consists in studying the flow roots of “small” graphs. By computing the flow roots of small graphs with high girth (up to 32 vertices and girth at least 7),³ Haggard, Pearce, and Royle [22] have very recently found an explicit counterexample to the Welsh conjecture: the flow polynomial of the generalised Petersen graph $G(16, 6)$ has two real roots larger than $Q = 4$: $Q_1 \approx 4.0252205$, and $Q_2 \approx 4.2331455$. However, the same authors conjectured the following modification of Conjecture 1.3, in which 4 is replaced by 5, and the endpoint $Q = 5$ is now included in accordance with Tutte’s five-flow conjecture:

¹Obviously, $\Phi_G(Q) = 0$ if G has a bridge, because of the “nowhere zero” condition.

² Indeed, the Petersen graph—which is a special case $G(5, 2)$ of the generalised Petersen graphs $G(n, k)$ to be defined in section 3 below—has the flow polynomial $\Phi_{G(5,2)}(Q) = (Q-1)(Q-2)(Q-3)(Q-4)(Q^2-5Q+10)$, which vanishes at $Q = 4$.

³Note that Kochol [35] proved that the smallest counterexample of Tutte’s five-flow conjecture should have girth at least 9.

Conjecture 1.6 (Haggard–Pearce–Royle [22]) *For any bridgeless graph G , $\Phi_G(Q) > 0$ for $Q \in [5, \infty)$.*

Even though the naive parallelism between $\chi_G(Q)$ and $\Phi_G(Q)$ has been invalidated by the above-mentioned counterexample to Conjecture 1.3, a related line of reasoning would be that all these conjectures and theorems might be related by replacing Q for the chromatic polynomial with $Q+1$ for the flow polynomial. Thus, the four-colour theorem [1] “translates” into the Tutte five-flow conjecture [53, 54]. Similarly, Conjecture 1.6 is the “translated” version of the Birkhoff–Lewis conjecture. (Note that the translation of Royle’s result [41] is consistent with the fact that Conjecture 1.3 [57] is false.)

In this paper we study the flow polynomial on the infinite family of graphs known as the generalised Petersen graphs. Our main results are the following:

Theorem 1.7 *The value $Q = 5$ is an isolated accumulation point of real zeros of the flow polynomial $\Phi_G(Q)$ for the families of bridgeless graphs $G(6n, 6)$ and $G(7n, 7)$ with $n \geq 3$. Moreover,*

- (a) *There is a sequence of real zeros $\{Q_n\}$ of the flow polynomials $\Phi_{G(6n, 6)}$ that converges to $Q = 5$ from below.*
- (b) *There is a sequence of real zeros $\{Q_n\}$ of the flow polynomials $\Phi_{G(7n, 7)}$ that converges to $Q = 5$. The sub-sequence with odd (resp. even) n converges to $Q = 5$ from above (resp. below).*

Theorem 1.8

- (a) *The bridgeless graph $G(119, 7)$ has flow roots at $Q \approx 5.00002$ and $Q \approx 5.16534$ (where $\approx$ means “within 10^{-5} ”).*
- (b) *The value $Q_c(7) \approx 5.235261$ (where $\approx$ means “within 10^{-6} ”) is an accumulation point of real zeros of the flow polynomials $\Phi_{G(7n, 7)}(Q)$. In particular, the sub-sequence for odd n of the real zeros $\{Q_n\}$ of the flow polynomials $\Phi_{G(7n, 7)}$ converges to $Q_c(7)$ from below.*

Remark. The largest real flow root we have explicitly found is $Q_0 \approx 5.1653424423$.

Thus, the Welsh conjecture and the “translated Birkhoff–Lewis conjecture” are false, and the Tutte conjecture is “almost false” (in the same sense that the four-colour theorem is “almost false” [8, 41]). On the other hand, Theorem 1.7 includes the “translated version” of that of Royle [41] for real chromatic roots.

In this work, we have considered the family of graphs $G(nk, k)$ with $k \leq 7$ and $n > 2$. For each k , we have located the set of accumulation points in the complex Q -plane of the roots of the flow polynomial $\Phi_{G(nk, k)}$, as $n \rightarrow \infty$. Most accumulation points belong to limiting curves $\mathcal{B}_k$; and in particular, we are interested in locating the points $Q_c(k)$, defined as the largest real value where the limiting curves $\mathcal{B}_k$ cross the real axis. (These points are likely to be accumulation points as $n \rightarrow \infty$ of real zeros, as in Theorem 1.8(b); but not always: see section 6.3.) We have been able to obtain the values of $Q_c(k)$ for $k \leq 11$; and the numerical

extrapolation of these values to $k \rightarrow \infty$ yields $\lim_{k \rightarrow \infty} Q_c(k) = Q_0 \approx 5.75$. We expect that this value is the largest real accumulation point one can get from the family $G(nk, k)$.

Based on this—and on the failure of Conjectures 1.3 and 1.6—we venture the following weaker conjecture:

Conjecture 1.9 *For any bridgeless graph G , $\Phi_G(Q) > 0$ for $Q \in [6, \infty)$.*

This translated version of the Birkhoff–Lewis theorem [15] bears the same relation to the Seymour theorem [46] as the original Birkhoff–Lewis theorem bears to the 5-colour theorem, or the Birkhoff–Lewis conjecture does to the 4-colour theorem; this can be taken as (weak) evidence that Conjecture 1.9 might be true. On the other hand, firm believers of the translation mechanism outlined above would conclude—by reverse translation of our Theorem 1.8 (and its extension)—that the Birkhoff–Lewis conjecture is false! We do not, however, possess any evidence whatsoever in support of this conclusion, based on actual studies of the chromatic polynomial.

Note that we do in fact exhibit *infinitely* many flow roots in an interval $Q \in [4, Q_0]$ with $Q_0 > 5$. This means, loosely speaking, that one should look for a purely combinatorial proof of Tutte’s 5-flow conjecture. This is exactly the same situation as for the 4- and 5-colour theorems; they hold true even though one can find families of graphs with real roots approaching $Q = 4$ from below [41], and other families with complex roots approaching densely to $Q = 4$ and $Q = 5$ [48]. If Conjecture 1.9 turns out to be false, then it is very plausible that there exist *no upper bound* for real flow roots of arbitrary bridgeless graphs.

On a more technical level, we exhibit a method for computing exactly the flow polynomial on very large generalised Petersen graphs (which can readily be adapted to other similar graph families). This method relies on a transfer matrix construction similar to the one employed in our previous work [26, 27] on the chromatic polynomial for graphs with periodic longitudinal boundary conditions.

The paper is organised as follows. In section 2 we define the flow polynomial carefully and exhibit its relation to the Q -state Potts model. Building on this, we show in section 3 how the flow polynomial on generalised Petersen graphs can be built by a transfer matrix construction. Our results, given in section 4, are obtained by implementing this construction on a computer and pushing the computation to as large graphs as possible. Note that although obtained by computational means, the flow polynomials are exact and involve no approximation whatsoever. In section 5 we introduce the Beraha–Kahane–Weiss theorem, that plays an important role in establishing our results. In section 6, we describe our analytic findings about the real zeros of the flow polynomial for this family of graphs, and in section 7, we summarise the more experimental (i.e., numerical) aspects of our work. To conclude, section 8 contains a discussion of our results. In Appendix A, we prove some technical lemmas included in the text, and in Appendix B, we give the coefficients of the generalised Petersen graph $P(119, 7)$.

2 Flow polynomial

Let $G = (V, E)$ be a connected graph and Γ be an Abelian group. Assign an arbitrary orientation to each edge $e \in E$. With respect to any fixed vertex $i \in V$, the edges E_i incident on i can then be characterised as either ingoing or outgoing: $E_i = E_i^{\text{in}} \cup E_i^{\text{out}}$.

A Γ -flow on G is a map $\phi: E \mapsto \Gamma$ that attributes a variable $\phi(e)$ to each edge $e \in E$, subject to the constraint

$$\sum_{e \in E_i^{\text{in}}} \phi(e) = \sum_{e \in E_i^{\text{out}}} \phi(e) \quad (2.1)$$

for any $i \in V$. The edge orientation is actually immaterial in these definitions: if one wants to change the orientation of an edge e_0 , it suffices to change simultaneously the sign of the flow along that edge, $\phi(e_0) \rightarrow -\phi(e_0)$.

A *nowhere zero* Γ -flow is a Γ -flow ϕ such that $\phi(e) \neq 0$ for all $e \in E$. If Γ is a *finite* Abelian group of order Q , then the number of nowhere zero Γ -flows depends only on Q , and is called the flow polynomial $\Phi_G(Q)$, for reasons we shall exhibit in the next paragraph. In particular, a $\mathbb{Z}_Q$ -flow (resp. a *nowhere zero* $\mathbb{Z}_Q$ -flow) on G is a map $\phi: E \mapsto \{0, 1, \dots, Q-1\}$ (resp. $\phi: E \mapsto \{1, 2, \dots, Q-1\}$) for which the constraint (2.1) is imposed modulo Q .

Clearly, the total number of $\mathbb{Z}_Q$ -flows on G is $Q^{c(E)}$, where for any subset $E' \subseteq E$, $c(E')$ denotes the number of independent cycles (cyclomatic number) in the induced graph $G' = (V, E')$. To obtain the number of nowhere zero $\mathbb{Z}_Q$ -flows, we first subtract for each $e \in E$ the flows for which $\phi(e) = 0$. Since flows with two zero-flow edges will be subtracted off twice, these must be put back in the sum, and proceeding by inclusion-exclusion we find [61]

$$\Phi_G(Q) = \sum_{E' \subseteq E} (-1)^{|E| - |E'|} Q^{c(E')} . \quad (2.2)$$

By this result, $\Phi_G(Q)$ is indeed a polynomial in Q , and (2.2) is used to extend the definition to $Q \in \mathbb{C}$.

Meanwhile, recall the partition function of the Q -state Potts model [39]

$$Z_G(Q, v) = \sum_{\sigma} \prod_{(ij) \in E} e^{K \delta(\sigma(i), \sigma(j))} , \quad (2.3)$$

where the map $\sigma: V \rightarrow \{0, 1, \dots, Q-1\}$ is called the spin, and K is the coupling constant. The Kronecker delta function $\delta(x, y)$ is defined by $\delta(x, y) = 1$ if $x = y$, and $\delta(x, y) = 0$ otherwise. We have introduced the convenient parameter $v = e^K - 1$. By expanding the edge product and performing the sum over σ , one recovers the partition function in the Fortuin-Kasteleyn cluster representation [20]

$$Z_G(Q, v) = \sum_{E' \subseteq E} v^{|E'|} Q^{k(E')} , \quad (2.4)$$

where $k(E')$ is the number of connected components in $G' = (V, E')$.

Graph theorists will recognise in (2.4) [a reparametrisation of] the Tutte polynomial [53] and interpret σ as a vertex colouring. Proper vertex colourings, i.e., those for which adjacent vertices are coloured differently, are obtained for $K \rightarrow -\infty$, and therefore

$$\chi_G(Q) = Z_G(Q, -1) \quad (2.5)$$

is the chromatic polynomial.

Setting instead $v = -Q$ in (2.4), and using the topological identity

$$k(E') = |V| - |E'| + c(E') , \quad (2.6)$$

one establishes the connection with the flow polynomial

$$\Phi_G(Q) = (-1)^{|E|} Q^{-|V|} Z_G(Q, -Q). \quad (2.7)$$

In the case where G is planar, let G^* denote the dual graph. Recall the fundamental duality relation [62] of the Potts model partition function

$$Z_G(Q, v) = K Z_{G^*}(Q, v^*), \quad (2.8)$$

where v^* is the dual of v

$$v v^* = Q, \quad (2.9)$$

and the proportionality factor is

$$K = Q^{1-|V^*|} v^{|E|} = Q^{|V|-|E|-1} v^{|E|}. \quad (2.10)$$

Noticing that $v = -Q$ is dual to $v = -1$ by (2.9) furnishes a relation between the flow polynomial of G and the chromatic polynomial of G^* . Indeed, using (2.8) and (2.10) we have [51]

$$\chi_{G^*}(Q) = Q \Phi_G(Q). \quad (2.11)$$

Note that $\Phi_G(Q) = 0$ if G contains a bridge $e_0 \in E$. Indeed, by the constraint (2.1) one would have $\phi(e_0) = 0$, preventing the existence of a nowhere zero flow. This observation is consistent with the duality relation (2.11), since a bridge in G implies a loop in G^* .

Alternatively, the relation (2.11) can be proved by noting that there exists an obvious bijection between the nowhere zero $\mathbb{Z}_Q$ -flows on G and the proper colourings of the faces of G , with the colour on one face being fixed. Indeed, let ϕ be a flow on G . Then, turning around a vertex, each time one moves from a face i to an adjacent face j , if the separating edge e is seen oriented to the right (resp. left), its flow variable $\phi(e)$ defines the colour difference $c_j - c_i = \phi(e)$ (resp. $c_j - c_i = -\phi(e)$). Starting from the face with fixed colour, these differences define the face colouring of the whole graph. The mapping from proper colourings to flows follows similarly.

It is useful to note that for any bridgeless 3-connected graph G , one can deduce from (2.4)/(2.7) that $\Phi_G(Q)$ is a polynomial in Q of degree $|E| - |V| + 1$ in Q , and that the first two coefficients of $\Phi_G(Q)$ are given by

$$\Phi_G(Q) = Q^{|E|-|V|+1} - |E| Q^{|E|-|V|} + \dots \quad (2.12)$$

The first term comes from the fact that there is a unique spanning graph (V, E') in (2.4)/(2.7) with $E' = E$. The second term is given by the contribution of the $|E|$ spanning subgraphs (V, E') with $E' = E \setminus e$ for each $e \in E$, and the observation that $k(E') = 1$ since G is connected and bridgeless. Notice that if we consider the spanning graph (V, E') with $E' \setminus \{e, e'\}$ for any two distinct edges $e, e' \in E$, the 3-connectedness of G guarantees that the next term in (2.12) will be of order $Q^{|E|-|V|-1}$, as in this case we also have $k(E') = 1$.

3 Transfer matrix for flow polynomials of generalised Petersen graphs

3.1 Generalised Petersen graphs


The goal of this paper is to evaluate the flow polynomial on a family of graphs $G(n, k)$ called *generalised Petersen graphs* and defined as follows: Let n, k be positive integers such that $n > k$. Then $G(n, k)$ is a cubic graph with $2n$ vertices denoted i_p and j_p for $p = 1, 2, \dots, n$: i.e.,

$$V(G(n, k)) = \{i_1, \dots, i_n, j_1, \dots, j_n\}. \quad (3.1)$$

The edge set consists of $3n$ edges $(i_p j_p)$, $(i_p i_{p+1})$, $(j_p j_{p+k})$, for $p = 1, 2, \dots, n$, and with all indices considered modulo n : i.e.,

$$E(G(n, k)) = \{(i_p, j_p), (i_p i_{p+1}), (j_p j_{p+k}) \mid 1 \leq p \leq n\}. \quad (3.2)$$

Note that $G(n, k)$ is simple for $n \neq 2k$; but it has double edges when $n = 2k$. These graphs were introduced by Watkins [56]. As an example, $G(12, 4)$ can be drawn as follows:



The graphs $G(n, k)$ are clearly bridgeless. They are non-planar for all pairs (n, k) , except for the case $(3, 2)$, and the two sub-families $(p, 1)$ and $(2p, 2)$ with $p \geq 1$. They have girth 8 for n and k sufficiently large. We have thus a two-parameter family of non-planar cubic graphs with high girth, and based on exhaustive studies of small graphs [22] we expect this family to produce large real flow roots.

Remark. Note that we do not expect that any of these roots to be $Q = 5$. In fact, the generalised Petersen graphs satisfy the following

Lemma 3.1 *For every generalised Petersen graph $G(n, k)$ with n, k positive integers such that $k < n$, $\Phi_{G(n, k)}(5) > 0$. In fact, every graph $G(n, k)$ other than the ordinary $G(5, 2)$ has $\Phi_{G(n, k)}(4) > 0$.*

PROOF. It is well known [16] that every generalised Petersen graph $G(n, k)$ (with the exception of the Petersen graph $G(5, 2)$ itself), admits a Tait colouring: i.e., an edge 3-colouring

such that at every vertex, the three incident edges take distinct colours.⁴ The existence of such edge 3-colourings is equivalent, for cubic and loopless graphs, to the existence of a nowhere-zero 4-flow for the same graph [32, Proposition 2(b)]. Therefore, for all $G(n, k)$, except the Petersen graph $G(5, 2)$, $\Phi_{G(n, k)}(4) > 0$, and furthermore, $\Phi_{G(n, k)}(5) > 0$ by Proposition 1.1. The case $G(5, 2)$ is dealt with directly: from the exact expression for $\Phi_{G(5, 2)}$ (see footnote 2), we conclude that $\Phi_{G(5, 2)}(5) = 240 > 0$. ■

We shall however show that the five-flow conjecture is “almost false”, in the sense of Theorem 1.7.

3.2 Potts model transfer matrix

We wish to evaluate $Z_{G(n, k)}(Q, v)$ —of which the flow polynomial $\Phi_{G(n, k)}(Q)$ is a special case—by a transfer matrix construction.

Contrary to an often repeated but false statement, evaluating $Z_G(Q, v)$ by a transfer matrix construction is possible for *any* graph G , and does not require G to consist of a number of identical layers [7]. However, when G does have a layered structure—as is the case here— $Z_G(Q, v)$ can be computed by the repeated application of the *same* transfer matrix.

Supposing for simplicity that n is a multiple of k , the generalised Petersen graph $G(n, k)$ can be redrawn as in Figure 1. This turns $G(n, k)$ into a graph of n/k identical layers of width $L = k + 1$ vertices with periodic boundary conditions in the vertical direction. (We shall henceforth refer to this as periodic *longitudinal* boundary conditions, in accordance with the fact that the transfer matrix builds up the graph vertically.) We now claim that this implies that $Z_{G(n, k)}(Q, v)$ can be written as a Markov trace

$$Z_{G(n, k)}(Q, v) = \text{Tr}(\mathbb{T}_L)^{n/k} \quad (3.3)$$

of the (n/k) th power of a transfer matrix $\mathbb{T}_L$ to be defined shortly.

In general, for a layered graph of width L , $\mathbb{T}_L$ acts on basis states A_L which are set partitions of $2L$ points $\{1, 2, \dots, L, 1', 2', \dots, L'\}$. These basis states can be depicted as *partition diagrams*, which are hypergraphs on $2L$ vertices, drawn inside a rectangle with L vertices (labelled $1', 2', \dots, L'$) on top and L vertices (labelled $1, 2, \dots, L$) on bottom. Each hyperedge represents one block in the partition. A block that contains at least one vertex from both the top and bottom rows is called a *link*. A block containing precisely one vertex is called a *singleton*. The number of links in a diagram d is denoted $\ell(d)$.

⁴It is worth noting that the definition of the generalised Petersen graph $G(n, k)$ in Refs. [16, 56] explicitly excludes the case $n = 2k$. However, it is easy to see that any $G(2k, k)$ has a Tait colouring: e.g., the edges (i_p, j_p) take colour 1, the edges (i_p, i_{p+1}) take alternatively colours 2 and 3 (as p goes from 1 to n), and for each p , one of the double edges (j_p, j_{p+k}) takes colour 2, and the other edge, colour 3.





Figure 1: (a) Generalised Petersen graph $G(n, k)$, here with $k = 4$. There are n layers of two vertices in the vertical direction, but some edges link layers at distance k . The boundary conditions are periodic in the vertical direction. (b) When $n \in k\mathbb{N}$, $G(n, k)$ can be redrawn as shown. There are n/k layers of width $k + 1$ vertices, each comprising a total of $2k$ vertices. All edges now link vertices within the same layer, or in two adjacent layers.


The following example with $L = 4$



represents the partition $(12'3')(231'4')(4)$. It has two links and one singleton.

The multiplication $d = d_2 \cdot d_1$ of two partition diagrams is defined by stacking the diagrams vertically. Specifically, the top row of d_2 becomes the top row of d , the bottom row of d_1 becomes the bottom row of d , and the top row of d_1 is identified with the bottom row of d_2 . Any blocks not containing points in the top or bottom rows of d are removed in the process.

This gives, for example:



This diagram multiplication turns A_L into an associative *partition monoid* [23, 36] with identity $I = (11')(22') \cdots (LL')$. Observe that

$$\ell(d_2 \cdot d_1) \leq \min(\ell(d_2), \ell(d_1)). \quad (3.4)$$


The idea is now that these diagrams will represent the edge subset appearing in the cluster representation (2.4) of the Potts model partition function. The factors of v can be dealt with locally, and the tricky part is to get a handle on the non-local factors Q . To this end, it is natural to associate an element of $\mathbb{C}$ with each diagram, which will play the role of the Boltzmann weight, i.e., the weight of a partially built configuration E' in (2.4). In the diagram multiplication $d = d_2 \cdot d_1$, let $\kappa(d_1, d_2)$ be the number of blocks which are removed because they contain no point in the top or bottom rows of d . The non-local part of the Boltzmann weight is then $Q^{\kappa(d_1, d_2)}$.

These considerations motivate the definition of the *partition algebra* [23, 36] $\mathbb{C}A_L(Q)$ as the associative algebra over $\mathbb{C}$ with basis A_L and multiplication defined by

$$d_2 d_1 = Q^{\kappa(d_1, d_2)} (d_2 \cdot d_1). \quad (3.5)$$

The partition algebra $\mathbb{C}A_L(Q)$ can be represented faithfully as an algebra of matrices whose rows and columns are indexed by the partition monoid A_L . This is indeed the point of view that we shall take when constructing the transfer matrix of the flow polynomial and manipulating it explicitly (see section 4.2). The elements of A_L can then be interpreted as the basis states of this representation. With no risk of confusion, we shall therefore use the notation A_L to refer both to the partition algebra and to the set of basis states.

We now define a set of generators for the monoid A_L . These generators will be the elementary building blocks used to define the transfer matrix $\mathbb{T}_L$. Apart from the identity I , the necessary generators are the *join operators* J_{ij} that amalgamate the blocks containing points i, j, i' and j' , and the *detach operators* D_i that remove point i' from its block and turn it into a singleton. In the pictorial representation this gives rise to the diagrams



As a consequence of the above definitions, the product $D_i d$ within $\mathbb{C}A_L(Q)$ produces a factor Q if the point i' is a singleton in diagram d , and a factor 1 otherwise. In particular, we have

$$D_i^2 = Q D_i. \quad (3.6)$$

From these building blocks we can now form the operators representing the addition of an edge to the graph that is being built up by the transfer matrix. These are

$$\mathbf{H}_{ij} = I + v \mathbf{J}_{ij}, \quad (3.7a)$$

$$\mathbf{V}_i = vI + \mathbf{D}_i. \quad (3.7b)$$

The letters H and V stand for horizontal and vertical, where a horizontal edge is understood to link vertices within the same layer of the graph (recall Figure 1), and a vertical edge links vertices from two adjacent layers.

Inspecting Figure 1, and labelling the points as in the figure, we can now finally define the transfer matrix (with $L = k + 1$):

$$\mathbf{T}_L = \mathbf{H}_{01} \left(\prod_{i=k}^2 \mathbf{V}_0 \mathbf{H}_{0i} \right) \left(\prod_{i=0}^k \mathbf{V}_i \right) \quad (3.8a)$$

$$= \mathbf{H}_{01} \mathbf{V}_0 \mathbf{H}_{02} \mathbf{V}_0 \mathbf{H}_{03} \cdots \mathbf{V}_0 \mathbf{H}_{0k} \mathbf{V}_k \mathbf{V}_{k-1} \cdots \mathbf{V}_0 \quad (3.8b)$$

Note the order of indices in the products.

3.3 Markov trace and eigenvalue amplitudes

It remains to explain the meaning of the Markov trace Tr in (3.3). The Markov trace of any partition diagram $d \in A_L$ is by definition $Q^{\kappa(d)}$, where $\kappa(d)$ is the number of connected components in the diagram obtained from d by identifying the points i and i' for all $i = 1, 2, \dots, L$. This identification corresponds to implementing the periodic longitudinal boundary conditions in Figure 1. The definition of the Markov trace extends to the partition algebra $\mathbb{C}A_L(Q)$ by linearity.

With this definition, (3.3), and (3.8), we are in principle equipped to compute the partition function $Z_{G(n,k)}(Q, v)$ as a polynomial in Q and v . A practical problem for going to large k is however that the dimension of $\mathbf{T}_L$, i.e., the number of basis states A_L , grows very fast with L :

$$\dim \mathbf{T}_L = |A_L| = B_{2L}, \quad (3.9)$$

where B_n are the Bell numbers with exponential generating function (egf)

$$\sum_{n=0}^{\infty} \frac{B_n z^n}{n!} = \exp(e^z - 1). \quad (3.10)$$

Considerable progress can nevertheless be made if one takes advantage of the structure of the partition algebra [23]. In practical terms this means that the number of points participating in the partitions can be halved from $2L$ to L . We now explain how this comes about.

Denote by $A_L^{(\ell)}$ the elements of the partition monoid A_L with exactly ℓ links, and define for $i = 0, 1, \dots, L$

$$\mathcal{A}_{L,i} = \bigcup_{\ell=0}^i A_L^{(\ell)}. \quad (3.11)$$

Thanks to (3.4), the $\mathcal{A}_{L,i}$ are in fact ideals which, moreover, constitute a filtration of the monoid:

$$\mathcal{A}_{L,0} \subseteq \mathcal{A}_{L,1} \subseteq \cdots \subseteq \mathcal{A}_{L,L} = A_L. \quad (3.12)$$

This implies immediately that any operator $\mathbf{O}_L$ (and $\mathbf{T}_L$ in particular) in the partition algebra $\mathbb{C}A_L(Q)$ has a block-triangular structure with respect to ℓ . The eigenvalues of $\mathbf{O}_L$ can therefore be found by restricting to $A_L^{(\ell)}$. In this restriction, the $\mathbb{C}$ -weight of an operator that acts on $\mathbb{C}A_L^{(\ell)}(Q)$ by diminishing ℓ will be set to zero. From the point of view of the matrix representation of $\mathbb{C}A_L(Q)$, this restriction amounts to replacing a matrix $\mathcal{O}_L$ by another matrix $\mathcal{O}'_L$ in which all the off-diagonal blocks have been set to zero. Although $\mathcal{O}'_L$ does not belong to $\mathbb{C}A_L(Q)$, it is still a well-defined matrix and we can study its eigenvalues, which are the same as those of $\mathcal{O}_L$.

Now ℓ is conserved, and $\mathbf{O}_L$ cannot change the blocks of the partition containing *only* points from the bottom set $\{1, 2, \dots, L\}$, since the multiplication has been defined by acting on the top points only. Therefore, in the restriction to $A_L^{(\ell)}$, $\mathbf{O}_L$ is block-diagonal, with a number of identical blocks on the diagonal that corresponds to the number of partitions of the bottom points that do not form part of a link.

As far as the determination of the eigenvalues goes, one can therefore restrict further the basis states of $A_L^{(\ell)}$ to partitions of the top points $\{1', 2', \dots, L'\}$ only, with precisely ℓ marked blocks (which were the links in the full partition monoid). Note that the marked blocks are *distinguishable*, since the action of $\mathbf{O}_L$ can still exchange their order (relative to the now forgotten fixed order of the links with respect to the bottom points $\{1, 2, \dots, L\}$). One can then finally block-diagonalise $\mathbf{O}_L$ by rearranging these restricted basis states into linear combinations that are irreducible representations λ of the symmetric group S_ℓ .

To summarise, all distinct eigenvalues of $\mathbf{O}_L$ can be found by studying the irreducible representations labelled by ℓ and λ . We have thus the following decomposition of the Markov trace

$$\text{Tr } \mathbf{O}_L = \sum_{\ell=0}^L \sum_{\lambda \in S_\ell} \alpha_{\ell,\lambda} \text{tr}_{\ell,\lambda} \mathbf{O}_L, \quad (3.13)$$

where now $\text{tr}_{\ell,\lambda}$ are ordinary matrix traces. The coefficients $\alpha_{\ell,\lambda}$ are eigenvalue amplitudes, which can also be interpreted as the dimensions of the commutant of the partition algebra.

Consider now $\lambda \in S_\ell$ through its corresponding Young diagram, $Y(\lambda) = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$, where λ_i is the number of boxes in the i 'th row. If there is less than ℓ rows in $Y(\lambda)$ the expression is of course padded with zeros. One then has the result [23, Proposition 3.24]

$$\alpha_{\ell,\lambda} = \frac{\dim \lambda}{\ell!} \prod_{i=0}^{\ell-1} (Q - i - \lambda_{\ell-i}). \quad (3.14)$$

We recall that the dimension $\dim \lambda$ of the representation λ is given by the hook formula [42]

$$\dim \lambda = \frac{\ell!}{\prod_{x \in Y(\lambda)} h_x}, \quad (3.15)$$

where h_x is the hook length of the box $x \in Y(\lambda)$, i.e., the number of boxes to its right, plus the number of boxes below it, plus the box itself. We shall sometimes need the total

amplitude β_ℓ for a given number of marked blocks. This reads then

$$\beta_\ell = \sum_{\lambda \in S_\ell} \alpha_{\ell, \lambda} \dim \lambda. \quad (3.16)$$

Note that the non-planarity of the graph is crucial for the applicability of (3.13)–(3.14). Indeed, had the graph been planar and with periodic *transverse* boundary conditions (in addition to the periodic longitudinal boundary conditions that we assume throughout), the transfer matrix could only have changed the cyclic order of the links, and the relevant group would not have been S_ℓ , but rather the cyclic group C_ℓ . Its representation theory leads to very different expressions [40] for (3.13)–(3.14). A similar remark holds for planar graphs with free transverse boundary conditions, in which case the links cannot be permuted at all, and only the decomposition according to the value of ℓ is relevant.

3.4 Flow polynomial transfer matrix

The transfer matrix that produces the flow polynomial $\Phi_{G(n,k)}(Q)$ can be taken simply as T_L of sections 3.2–3.3, i.e., by specialising (3.3) to $v = -Q$.

One can however diminish the dimension of the relevant partition algebra by remarking that for $v = -Q$, the vertical operator V_i in (3.7b) is a projector (up to a constant). Indeed, by (3.6) one finds $\mathsf{V}_i^2 = (vI + \mathsf{D}_i)^2 = v^2I + (2v + Q)\mathsf{D}_i$, which is a multiple of V_i if and only if $v = -Q$. The normalised projector $(-Q)^{-1}\mathsf{V}_i$ annihilates any partition diagram in which the point i' is a singleton. Concerning the reduced partitions of the points $\{1', 2', \dots, L'\}$ with precisely ℓ marked blocks—as described in section 3.3—the precise statement is: $(-Q)^{-1}\mathsf{V}_i$ annihilates any reduced diagram in which i' is an un-marked singleton.

Since V_i and V_j commute for any i, j , the following operator

$$\mathsf{P}_L = (-Q)^{-L} \prod_{i=0}^{L-1} \mathsf{V}_i \quad (3.17)$$

is also a projector. It annihilates any reduced diagram containing an un-marked singleton. We can therefore replace (3.3) by

$$Z_{G(n,k)}(Q, -Q) = \text{Tr}(\tilde{\mathsf{T}}_L)^{n/k}, \quad (3.18)$$

where $\tilde{\mathsf{T}}_L = \mathsf{P}_L \mathsf{T}_L$ with $v = -Q$, and consider the trace only over states without un-marked singletons. This implies that the flow polynomial [cf. (2.7)] can be finally written as

$$\Phi_{G(n,k)}(Q) = (-1)^n Q^{-2n} \text{Tr}(\tilde{\mathsf{T}}_L)^{n/k}, \quad (3.19)$$

as the generalised Petersen graph $G(n, k)$ has $3n$ edges and $2n$ vertices. The prefactor $(-1)^n Q^{-2n}$ can be absorbed in the definition of the transfer matrix: if we define

$$\hat{\mathsf{T}}_L = (-1)^k Q^{-2k} \tilde{\mathsf{T}}_L, \quad (3.20)$$

then (3.19) becomes

$$\Phi_{G(n,k)}(Q) = \text{Tr}(\hat{\mathsf{T}}_L)^{n/k}. \quad (3.21)$$

The decomposition (3.13)–(3.14) goes through as before, now only with the “no unmarked singleton” constraint imposed on the representations labelled by ℓ and λ .

Remarks. 1. All the entries in the matrix $\tilde{T}_L$ are polynomials in Q ; but this property does not hold in general for the matrix elements of $\hat{T}_L$. Some of them may contain terms with inverse powers of Q .⁵

2. The structural properties of $\tilde{T}_L$ and $\hat{T}_L$ are obviously the same.

3.5 Dimensions of representations

Let us first consider the number of partitions $A_{k+1}^{(\ell)}$ of $k+1$ points with ℓ marked and distinguishable blocks. It is given by

$$|A_k^{(\ell)}| = k! [z^k] ((e^z - 1)^\ell \exp(e^z - 1)) , \quad (3.22)$$

as is easily seen by elementary manipulations of the egf of the Bell numbers (the case $\ell = 0$). Using (3.10), and the fact that the Stirling numbers of the second kind $\left\{ \begin{smallmatrix} k \\ \ell \end{smallmatrix} \right\}$ (or Stirling subset numbers) [21] have the following egf [19]

$$S^\ell(z) = \frac{1}{\ell!} (e^z - 1)^\ell \quad (3.23a)$$

$$\left\{ \begin{smallmatrix} k \\ \ell \end{smallmatrix} \right\} = k! [z^k] S^\ell(z) \quad (3.23b)$$

we can derive the following closed form for $|A_k^{(\ell)}|$:

$$|A_k^{(\ell)}| = \ell! \sum_{p=0}^k \binom{k}{p} \left\{ \begin{smallmatrix} p \\ \ell \end{smallmatrix} \right\} B_{k-p} \quad (3.24a)$$

$$= \ell! \sum_{s=0}^{k-\ell} \binom{\ell+s}{\ell} \left\{ \begin{smallmatrix} k \\ \ell+s \end{smallmatrix} \right\} \quad (3.24b)$$

where we have gone from (3.24a) to (3.24b) by using the well-known expression of the Bell numbers in terms of the Stirling subset numbers [19]

$$B_n = \sum_{s=0}^n \left\{ \begin{smallmatrix} n \\ s \end{smallmatrix} \right\} . \quad (3.25)$$

It is clear from (3.24) that $\ell! \mid |A_k^{(\ell)}|$, $|A_k^{(k)}| = k!$, and $|A_k^{(0)}| = B_k$.

Meanwhile, the sum of the eigenvalue amplitudes for a given ℓ and all possible Young diagrams λ is given by (3.14)–(3.16):

$$\beta_\ell = \ell! \sum_{i=0}^{\ell} \frac{(-1)^i}{i!} \binom{Q}{\ell-i} . \quad (3.26)$$

⁵Let us give an example for $k = 3$. When we apply the transfer matrix $\tilde{T}_4$ to the partition $(0'1'2'3')(0123)$, we get several partitions with coefficients that are polynomial in Q . In particular, we obtain the partition $(0'3')(1'2')(0123)$ with the coefficient $(-Q)^5$. If we divide this polynomial by the prefactor $(-1)^k Q^{-2k}$ [cf. (3.20)], we obtain $1/Q$, which is *not* a polynomial in Q .

This is indeed a polynomial in Q ; it can be rewritten in terms of the falling factorials $Q^i = \prod_{j=1}^i (Q + 1 - j)$ [21] as:

$$\beta_\ell = \sum_{i=0}^{\ell} (-1)^{\ell-i} \binom{\ell}{i} Q^i. \quad (3.27)$$

The values we need in this paper are:

$$\beta_0 = 1 \quad (3.28a)$$

$$\beta_1 = Q - 1 \quad (3.28b)$$

$$\beta_2 = Q^2 - 3Q + 1 \quad (3.28c)$$

$$\beta_3 = Q^3 - 6Q^2 + 8Q - 1 \quad (3.28d)$$

$$\beta_4 = Q^4 - 10Q^3 + 29Q^2 - 24Q + 1 \quad (3.28e)$$

$$\beta_5 = Q^5 - 15Q^4 + 75Q^3 - 145Q^2 + 89Q - 1 \quad (3.28f)$$

$$\beta_6 = Q^6 - 21Q^5 + 160Q^4 - 545Q^3 + 814Q^2 - 415Q + 1 \quad (3.28g)$$

$$\beta_7 = Q^7 - 28Q^6 + 301Q^5 - 1575Q^4 + 4179Q^3 - 5243Q^2 + 2372Q - 1 \quad (3.28h)$$

Just as in the case of the planar partition algebra [see Ref. [31], in particular Eqs. (2.16) and (2.20)] the compatibility between the dimensions $|A_k^{(\ell)}|$ and the amplitudes β_ℓ can be expressed in the form of a sumrule:

$$\sum_{\ell=0}^k \frac{1}{\ell!} \beta_\ell |A_k^{(\ell)}| = Q^k. \quad (3.29)$$

This expresses that the number of degrees of freedom per vertex of the graph is indeed Q , as expected. We also have the sumrule

$$B_k^{(0)} = \sum_{\ell=0}^k \frac{1}{\ell!} |A_k^{(\ell)}| = \sum_{r=0}^k 2^r \left\{ \begin{matrix} k \\ r \end{matrix} \right\}, \quad (3.30)$$

where the integers $B_n^{(0)}$ form the sequence A001861 of [47]. Their egf is

$$\sum_{n=0}^{\infty} \frac{B_n^{(0)} z^n}{n!} = \exp(2(e^z - 1)), \quad (3.31)$$

as can be deduced from (3.22).

For fixed k and ℓ , we now introduce the “no un-marked singletons” constraint. We then obtain a smaller set of partitions $\tilde{A}_{k+1}^{(\ell)}$ of cardinality $\tilde{N}_k(\ell) = |\tilde{A}_{k+1}^{(\ell)}|$.

The number of partitions of k points with ℓ marked and distinguishable blocks satisfying the “no un-marked singletons” constraint is

$$|\tilde{A}_k^{(\ell)}| = k! [z^k] ((e^z - 1)^\ell \exp(e^z - 1 - z)). \quad (3.32)$$

The key point is that the number of partitions with no singletons is given by the egf $\exp(e^z - 1 - z)$ [19, p. 111]. The numbers associated with this egf are given by

$$S_n = n! [z^n] \exp(e^z - 1 - z) = (-1)^n \sum_{q=0}^n \binom{n}{q} (-1)^q B_q, \quad (3.33)$$

where the B_n are the Bell numbers (3.10). Then, a closed form for the numbers $|\tilde{A}_k^{(\ell)}|$ reads:

$$|\tilde{A}_k^{(\ell)}| = \ell! \sum_{p=0}^k \binom{k}{p} \left\{ p \atop \ell \right\} S_{k-p}. \quad (3.34)$$

This formula implies that $\ell! \mid |\tilde{A}_k^{(\ell)}|$, and that $|\tilde{A}_k^{(k)}| = k!$.

The sumrule corresponding to (3.29) now reads

$$\sum_{\ell=0}^k \frac{1}{\ell!} \beta_\ell |\tilde{A}_k^{(\ell)}| = (Q-1)^k. \quad (3.35)$$

The fact that Q has been replaced by $Q-1$ is a manifestation of the “nowhere zero” constraint. We also have the sumrule corresponding to (3.30)

$$\tilde{B}_k^{(0)} = \sum_{\ell=0}^k \frac{1}{\ell!} |\tilde{A}_k^{(\ell)}| = \sum_{r=0}^k \binom{k}{r} B_r^{(0)} (-1)^{k-r}, \quad (3.36)$$

where the integers $B_n^{(0)}$ are given by (3.30). Their egf is

$$\sum_{n=0}^{\infty} \frac{\tilde{B}_n^{(0)} z^n}{n!} = \exp(2(e^z - 1) - z). \quad (3.37)$$

Notice that the difference $|A_k^{(\ell)}| - |\tilde{A}_k^{(\ell)}|$ gives the number of partitions of the set $\{1, 2, \dots, k\}$ with ℓ marked points and with at least one un-marked singleton. These partitions do not contribute to the final result, as they are associated to null eigenvalues.

4 Flow polynomial for the generalised Petersen graphs

4.1 General theory

Let us start with the simplest case $k=1$. The graph $G(n, 1)$ is isomorphic to a cyclic ladder of width 2. From the known Potts-model partition function [18, and references therein], one can easily derive

$$\Phi_{G(n,1)}(Q) = (Q^2 - 3Q + 1)(-1)^n + (Q-1)(Q-3)^n + (Q-2)^n, \quad (4.1)$$

where the eigenvalues $\mu = -1, Q-3, Q-2$ correspond to the sectors with $\ell = 2, 1, 0$ links, respectively.

Let us now focus on $k \geq 2$. The flow polynomial (3.21) for the generalised Petersen graph $G(nk, k)$ can be written as

$$\Phi_{G(nk,k)}(Q) = \sum_{\ell=0}^{k+1} \sum_{\lambda \in S_\ell} \alpha_{\ell,\lambda} \text{tr}_{\ell,\lambda}(\hat{\mathbf{T}}_{k+1})^n, \quad (4.2)$$

where the amplitudes $\alpha_{\ell,\lambda}$ are given by (3.14). This formula is the most general one. In terms of the non-zero eigenvalues $\mu_{k,\ell,\lambda,s}$ of the transfer matrix $\hat{T}_{k+1}$, it reads:

$$\Phi_{G(nk,k)}(Q) = \sum_{\ell=0}^{k+1} \sum_{\lambda \in S_\ell} \alpha_{\ell,\lambda} \sum_{s=1}^{\tilde{N}_k(\ell)} \mu_{k,\ell,\lambda,s}^n, \quad (4.3)$$

where $\tilde{N}_k(\ell) = |\tilde{A}_{k+1}^{(\ell)}|$ [cf. (3.32)/(3.34)].

The flow polynomial $\Phi_{G(nk,k)}$ (4.3) is obtained as a linear combination of ordinary matrix traces (i.e., sums of powers of the eigenvalues $\mu_{k,\ell,\lambda,s}$) with definite coefficients $\alpha_{\ell,\lambda}$ (3.14). This is *all* what we need to compute $\Phi_{G(nk,k)}$ rigorously from the various diagonal blocks of $\hat{T}_L$. It is worth stressing that Eq. (4.3) holds true irrespective of whether some eigenvalues happen to be identical or not. However, it will turn out useful in practice to have a “complete” decomposition of the flow polynomial, in the sense that there are no equalities among eigenvalues (except for what is accounted for in the multiplicities $\alpha_{\ell,\lambda}$). This reason motivates a deeper study of the eigenvalue structure of the transfer matrix $\hat{T}_L$.

To simplify the notation, we will denote $\tilde{T}_{k+1,\ell}$ the diagonal block of the full transfer matrix $\tilde{T}_{k+1}$ corresponding to partitions with exactly ℓ links. We will denote $\tilde{T}_{k+1,\ell,\lambda}$ the diagonal block of $\tilde{T}_{k+1,\ell}$ corresponding to the irreducible representation λ of the group S_ℓ . Similar notation will be used for the diagonals blocks of the transfer matrix $\hat{T}_{k+1}$.

For $1 \leq k \leq 7$, our exact computations (described in detail in section 4.2) have revealed the following structure described in Lemmas 4.1, and 4.2, Corollary 4.3, and Lemma 4.4 below:

Lemma 4.1 *Fix $1 \leq k \leq 7$. Then for any $0 \leq \ell \leq k+1$, the diagonal block $\hat{T}_{k+1,\ell}$ contains the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$ with multiplicity $\tilde{N}_{k,1}(\ell) \ell!$, where*

$$\tilde{N}_{k,1}(\ell) = \begin{cases} 0 & \text{if } \ell = 0 \\ \frac{1}{(\ell-1)!} |\tilde{A}_k^{(\ell-1)}| = \sum_{p=0}^k \binom{k}{p} \left\{ \begin{matrix} p \\ \ell-1 \end{matrix} \right\} S_{k-p} & \text{if } 1 \leq \ell \leq k+1 \end{cases} \quad (4.4)$$

and S_n is given in (3.33).

Remarks. 1. We have proven that this lemma can be extended to any $k \geq 1$. This general result corresponds to Lemma A.1 in Appendix A.

2. The eigenvalue $\mu_{k,k+1} = (-1)^k$ will be called “trivial” in the following, because it is directly related to the structural properties of the transfer matrix $\hat{T}_L$. In the proof of Lemma A.1, we show that, when we order the basis states appropriately, the diagonal block $\hat{T}_{k+1,\ell}$ has a block triangular structure with two diagonal blocks:

- One block is a diagonal matrix of dimensionality $\tilde{N}_{k,1}(\ell) \ell!$ with all diagonal entries equal to the trivial eigenvalue $\mu_{k,k+1}$.
- The other block is not in general a diagonal matrix, and gives rise to the “non-trivial” eigenvalues.

3. In particular, the general Lemma A.1 imply that for any $k \geq 1$ and $\ell = k + 1$, the multiplicity of the trivial eigenvalue $\mu_{k,k+1}$ is given by $\tilde{N}_k(k+1) = \tilde{N}_{k,1}(k+1)(k+1)! = (k+1)!$.

Lemma 4.2 *Fix $1 \leq k \leq 7$, and $\ell = k$. Then, there are k non-trivial eigenvalues $\mu_{k,k,s}$ in the diagonal block $\tilde{T}_{k+1,k}$ (once we have removed the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$). Each of these eigenvalues $\mu_{k,k,s}$ has multiplicity $k!$.*

Remarks. 1. We have proven that this lemma can be extended to any $k \geq 1$. This general result corresponds to Lemma A.2 in Appendix A.

2. For each irreducible representation λ of the group S_k , the diagonal block $\hat{T}_{k+1,k,\lambda}$ contains the same k non-trivial eigenvalues $\mu_{k,k,s}$.

Corollary 4.3 *Fix $1 \leq k \leq 7$. Then for all pairs (ℓ, λ) with $0 \leq \ell \leq k-1$ and $\lambda \in S_\ell$, the diagonal block $\tilde{T}_{k+1,\ell,\lambda}$ has $\tilde{N}_{k,0}(\ell)$ non-trivial eigenvalues $\mu_{k,\ell,\lambda,s}$ (once we have removed the trivial eigenvalue $\mu_{k+1,k} = (-1)^k$, which appears for every $1 \leq \ell \leq k+1$). The value of $\tilde{N}_{k,0}(\ell)$ is given by:*

$$\tilde{N}_{k,0}(\ell) = \begin{cases} S_{k+1} + \sum_{p=0}^k \binom{k+1}{p+1} S_{k-p} & \text{if } \ell = 0 \\ Y_\ell \sum_{p=1}^k \binom{k}{p} \left\{ \begin{matrix} p \\ \ell \end{matrix} \right\} [S_{k-p} + S_{k+1-p}] & \text{if } 1 \leq \ell \leq k-1 \\ k & \text{if } \ell = k \\ 0 & \text{if } \ell = k+1 \end{cases} \quad (4.5)$$

where S_n is given by (3.33), and Y_ℓ is the number of Young tableaux with ℓ cells.

Remarks. 1. The number Y_ℓ of Young tableaux with ℓ cells (sequence A000085 in [47]) has the following egf:

$$Y_\ell = \sum_{\lambda \in S_\ell} \dim \lambda = \ell! [z^\ell] \exp(z + z^2) \quad (4.6a)$$

$$Y_\ell = \sum_{p=0}^{\ell/2} \frac{(2p)!}{p! 2^p} \binom{\ell}{p} \quad (4.6b)$$

2. This corollary follows from Lemmas 4.1 and 4.2. As these two lemmas can be generalized for any $k \geq 1$, the more general Corollary A.3 in Appendix A follows.

Lemma 4.4 *Fix $1 \leq k \leq 7$. Then for each pair (ℓ, λ) with $0 \leq \ell \leq k$ and $\lambda \in S_\ell$, the non-trivial eigenvalues $\mu_{k,\ell,\lambda,s}$ (or $\mu_{k,k,s}$ when $\ell = k$) are all distinct and also different from the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$.*

Remarks. 1. We do not know how to prove the extension of this result to larger values of $k \geq 8$; but we conjecture that Lemma 4.4 holds true for any $k \geq 1$.

2. The fact that all eigenvalues $\mu_{k,\ell,\lambda,s}$ are distinct in the generic case, implies that we do not have to worry about the non-degenerate-dominance condition in the Beraha–Kahane–Weiss theorem (see section 5).

Lemmas 4.1, 4.2, and 4.4 show that for $1 \leq k \leq 7$, we have found that the decomposition (4.3) is “almost complete”, as all eigenvalues $\mu_{k,\ell,\lambda,s}$ are distinct, except for two cases:

- (a) The trivial eigenvalue $\mu_{k,k+1} = (-1)^k$, that appears in every sector with $1 \leq \ell \leq k+1$.
- (b) The eigenvalues $\mu_{k,k,s} = \mu_{k,k,\lambda,s}$ (of the sector $\ell = k$) do not depend on the value of the representation $\lambda \in S_k$.

This means that, if we want to have a “complete” decomposition of the flow polynomial, then the amplitudes for the eigenvalues $\mu_{k,k+1}$ and $\mu_{k,k,s}$ are not longer given by a single coefficient $\alpha_{\ell,\lambda}$; but by a linear combination of several coefficients. In particular, the coefficient associated to the eigenvalues $\mu_{k,k,s}$ should be equal to β_k (3.16)/(3.26). The coefficient γ_{k+1} associated to the trivial eigenvalue $\mu_{k,k+1}$ should be given by

$$\gamma_{k+1} = \beta_{k+1} + \sum_{\ell=1}^k \tilde{N}_{k,1}(\ell) \beta_{\ell}, \quad (4.7)$$

where $\tilde{N}_{k,1}(\ell)$ is given in (4.4).

Lemmas 4.1, 4.2, and 4.4, Corollary 4.3, and the above discussion imply that the flow polynomial for the generalised Petersen graph $G(nk, k)$ is given by the following “complete” decomposition:

Corollary 4.5 *Fix $1 \leq k \leq 7$. Then the flow polynomial for the generalised Petersen graph $G(nk, k)$ is given for any $n \geq 1$ by*

$$\Phi_{G(nk,k)}(Q) = \sum_{\ell=0}^{k-1} \sum_{\lambda \in S_{\ell}} \alpha_{\ell,\lambda} \sum_{s=1}^{\tilde{N}_{k,0}(\ell)} \mu_{k,\ell,\lambda,s}^n + \beta_k \sum_{s=1}^k \mu_{k,k,s}^n + \gamma_{k+1} (-1)^{nk} \quad (4.8)$$

where β_{ℓ} is given in (3.26)/(3.27), γ_{k+1} is given by (4.7), and all eigenvalues $\{\mu_{k,\ell,\lambda,s}, \mu_{k,k,s}, (-1)^k\}$ are distinct.

Remark. We conjecture that the above “complete” decomposition for the flow polynomial $\Phi_{G(nk,k)}$ holds true for any $k \geq 8$.

We can compute the coefficients γ_{k+1} by using (4.7), the expressions for the amplitudes β_{ℓ} (3.28), and the values of $\tilde{N}_{k,1}(\ell)$ (4.4). The results for $1 \leq k \leq 7$ are:

$$\gamma_2 = Q^2 - 3Q^2 + 1 \quad (4.9a)$$

$$\gamma_3 = Q^3 - 5Q^2 + 6Q - 1 \quad (4.9b)$$

$$\gamma_4 = Q^4 - 7Q^3 + 15Q^2 - 11Q + 1 \quad (4.9c)$$

$$\gamma_5 = Q^5 - 9Q^4 + 28Q^3 - 38Q^2 + 20Q - 1 \quad (4.9d)$$

$$\gamma_6 = Q^6 - 11Q^5 + 45Q^4 - 90Q^3 + 90Q^2 - 27Q + 1 \quad (4.9e)$$

$$\gamma_7 = Q^7 - 13Q^6 + 66Q^5 - 175Q^4 + 260Q^3 - 207Q^2 + 70Q - 1 \quad (4.9f)$$

$$\gamma_8 = Q^8 - 15Q^7 + 91Q^6 - 301Q^5 + 595Q^4 - 707Q^3 + 469Q^2 - 135Q + 1 \quad (4.9g)$$

For instance, for $k = 3$, we have that $\gamma_4 = \beta_1 + 4\beta_2 + 3\beta_3 + \beta_4$.

The total number of distinct eigenvalues as a function of k is given by

$$\tilde{D}_k = 1 + \sum_{\ell=0}^k \tilde{N}_{k,0}(\ell), \quad (4.10)$$

where the $\tilde{N}_{k,0}(\ell)$ are given in (4.5). The values are $\tilde{D}_k = 3, 7, 36, 229, 1658, 12803, 105934$ for $k = 1, \dots, 7$.

Remarks. 1. It is important to notice that, once the eigenvalues of the transfer matrix T_{k+1} [or those of $\hat{\mathsf{T}}_{k+1}$ (3.20)] are obtained, as well as their multiplicities, Corollary 4.5 establishes the *exact* form of the flow polynomial $\Phi_{G(nk,k)}$ for any $1 \leq k \leq 7$ and any $n \geq 1$. In particular, the exactness of the analysis does not depend on the validity of Lemma 4.4 for larger values of $k \geq 8$.

4.2 Practical procedure

We have written a PERL script to compute the *symbolic* transfer matrix T_{k+1} using ideas similar to those already explained in [26, 27]. For $1 \leq k \leq 4$, we have checked our programs using MATHEMATICA. Further checks were performed with code written in C that allows us to compute the leading eigenvalue for given values of k, ℓ , and $\lambda = (\ell)$ — the completely symmetric irreducible representation of S_ℓ .

The first step is to obtain the relevant diagonal blocks $\tilde{\mathsf{T}}_{k+1,\ell,\lambda}$ of the transfer matrix $\tilde{\mathsf{T}}_{k+1}$. We first fix the value of $0 \leq \ell \leq k+1$, and a bottom-row configuration compatible with the chosen value of ℓ and the “no un-marked singletons” condition. We then determine the basis of the relevant partition space of dimensionality $\tilde{N}_k(\ell)$ [cf. (3.34)]. Indeed, the result does not depend on the chosen bottom-row partition.⁶

We now choose an irreducible representation λ of the symmetric group S_ℓ of dimensionality $\dim \lambda$. To obtain the relevant diagonal block $\tilde{\mathsf{T}}_{k+1,\ell,\lambda}$ corresponding to λ , we simply take as our basis vector those linear combinations of the “standard” basis with the appropriate properties under S_ℓ . Finally, in order to “extract” the trivial eigenvalues⁷ Q^{2k} for $\ell \geq 1$, we exploit the structure of $\tilde{\mathsf{T}}_{k+1,\ell,\lambda}$ found in the proof of Lemma A.1.

In this way, we obtain all the diagonal blocks $\tilde{\mathsf{T}}_{k+1,\ell,\lambda}$ with $1 \leq k \leq 7$, $0 \leq \ell \leq k+1$, and all the $\lambda \in S_\ell$. We then have to compute the traces

$$\mathrm{tr} \hat{\mathsf{T}}_{k+1,\ell,\lambda}^n = \sum_{s=1}^{\tilde{N}_{k,0}(\ell)} \mu_{k,\ell,\lambda,s}^n, \quad (4.11)$$

for integer values of $n \geq 1$. For small values of k , this computation can be made symbolically for any n using a symbolic algebraic manipulator program, such as MATHEMATICA. However,

⁶To be more precise, there are bottom-row configurations that give rise to a basis of dimensionality $p\tilde{N}_k(\ell)$ [cf. (3.34)] for some integer $p \geq 2$. In these cases, one indeed obtains the same $\tilde{N}_k(\ell)$ eigenvalues; but each one repeated p times.

⁷Please note that the trivial eigenvalue for the matrix $\hat{\mathsf{T}}_{k+1}$ (3.20) is given by $\mu_{k,k+1} = (-1)^k$. But the corresponding eigenvalue for the matrix $\tilde{\mathsf{T}}_{k+1}$ is $(-1)^k(-1)^k Q^{2k} = Q^{2k}$.

for larger values of k (say, $k = 6, 7$) this is not feasible, as we have blocks of dimension as large as 11 816 (for $k = 7$, $\ell = 3$, and $\lambda = (2, 1)$).

A key issue in the subsequent analysis is the dependence of the above traces (4.11) on Q . The needed information is given by the following Lemma (the proof can be found in Appendix A):

Lemma 4.6 *Let $k \geq 1$, $0 \leq \ell \leq k + 1$, and λ be an irreducible representation of S_ℓ . Then $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ [cf. (4.11)] is a polynomial in Q of degree at most $n[k + \min(1 - \ell, 0)]$.*

We have made use of the following tricks:

- We know that the traces (4.11) are polynomials in Q of degree $d = n[k + \min(1 - \ell, 0)]$ by Lemma 4.6. Therefore, we only have to compute the *evaluation* of each trace at $d + 1$ integer values of $Q \neq 0$, and then reconstruct the corresponding polynomial using Lagrange's interpolation method. In order to check the result, we always compute at least $d + 2$ values of the trace. Please note that we compute the evaluation of the trace (4.11) by first computing the evaluation of the trace $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$, and then multiplying the result by the factor $(-1)^{nk} Q^{-2nk}$ [cf. (3.20)]. Indeed, this factor is not defined for $Q = 0$.
- *Not* all the entries of the transfer matrix $\tilde{T}_{k+1,\ell,\lambda}$ are integers for integer values of Q ; rather they are rational numbers. As we want to perform the trace computation with (infinite-precision) integer arithmetic, for each value of Q , we multiplied the matrix by the minimum (positive) integer value such that all entries are integers. After the computation is done, we reconstructed the true solution by dividing by the appropriate factor.
- The integers involved in the actual calculations are very large. Therefore, we compute the value of the trace for a given value of $Q \neq 0$ using modular arithmetic for a given set of prime numbers $p \leq 65521 \lesssim 2^{16}$ (we need up to 65 different primes). We then reconstruct the value using the Chinese remainder theorem using infinite-precision arithmetic in MATHEMATICA. We always use at least one more prime than needed, in order to check the result. To accelerate the computation of the trace using modular arithmetic, we use a program written in C.
- For $1 \leq k \leq 6$, we are able to compute the traces for many different values of n . However, for $k = 7$, the computation is so demanding, that we have focused on powers of the type $n = 2^q + 1$ with integer $q \geq 1$. The reason why we consider odd powers for $k = 7$ will become clear in section 6.

Once the traces are computed, we can form the flow polynomial using (4.8). Notice that for $1 \leq k \leq 7$, everything in this formula is exactly known. The zeros of the flow polynomials are obtained using the program MPSOLVE [12, 13]. This software has the advantage that if one requests the zeros with 50-digit precision (as in our case), the results are guaranteed to have at least such precision.

Remark. To give a clear idea of what has been achieved, consider the case $k = 7$, and more specifically the computation of Φ_G for the graph $G = G(17k, k)$, which is the largest

computation undertaken in this work. Applying naively (3.8) within the diagrammatic basis would imply computing the 17th power of T_{k+1} , which according to (3.9) is a matrix of dimension $B_{2(k+1)} = 10\,480\,142\,147$ whose entries are polynomials in Q of degree $2k$. The decomposition of T_{k+1} and use of the “no un-marked singleton” constraint has reduced the computation to the sum over 31 blocks, the largest of which has dimension 11 816. Even with these tricks, the computation took around six months, using 50–80 processors, corresponding to some 30 years of CPU time.

4.3 Additional checks

Because our results are derived using software, we have performed some tests in order to ensure that the results are right. First of all, for the smallest members of each family $G(nk, k)$, we have computed the flow polynomial using MAPLE. Indeed, both results agree in all cases.

For the cubic graphs $G(n, k)$ that we are considering, we may improve on (2.12) by adding a few more terms:

$$\begin{aligned} \Phi_{G(n,k)}(Q) &= Q^{|E|-|V|+1} - |E|Q^{|E|-|V|} + \left(\frac{|E|(|E|-1)}{2} - |V| \right) Q^{|E|-|V|-1} \\ &\quad - \left(\frac{|E|(|E|-1)(|E|-2)}{6} - |V|(|E|-2) \right) Q^{|E|-|V|-2} + \dots \end{aligned} \quad (4.12)$$

In this expression, the coefficient of $Q^{|E|-|V|-1}$ arises from two contributions in which $E \setminus E'$ is respectively two edges, and three edges all incident on the same vertex. The contributions to the coefficient of $Q^{|E|-|V|-2}$ are slightly more complicated to characterise. Inserting $|V| = 2n$ and $|E| = 3n$ we obtain

$$\Phi_{G(n,k)}(Q) = Q^{n+1} \left(1 - 3nQ^{-1} + \frac{n}{2}(9n-7)Q^{-2} - \frac{n}{2}(3n-2)(3n-5)Q^{-3} + \dots \right). \quad (4.13)$$

We have checked that for all the graphs $G(nk, k)$ we have considered, the flow polynomials obtained from the procedure outlined above indeed satisfy (4.13).

There are some theorems that give us some information about the location of the real zeros of the flow polynomial. The first theorem applies to a general bridgeless graph; while the second one applies only to cubic graphs (i.e., it is valid for $G(nk, k)$ with $n > 2$):

Theorem 4.7 (Wakelin [55]) *Let G be a bridgeless graph with $|V|$ vertices, $|E|$ edges, b blocks, and no isolated vertices. Then:*

- $\Phi_G(Q)$ is non-zero with sign $(-1)^{|E|-|V|+1}$ for $Q \in (-\infty, 1)$;
- $\Phi_G(Q)$ has a zero of multiplicity b at $Q = 1$.
- $\Phi_G(Q)$ is non-zero with sign $(-1)^{|E|-|V|+b+1}$ for $Q \in (1, \frac{32}{27}]$.

Theorem 4.8 (Jackson [24, 25]) *Let G be a 3-connected cubic graph with $|V|$ vertices and $|E|$ edges. Then:*

- $\Phi_G(Q)$ is non-zero with sign $(-1)^{|E|-|V|}$ for $Q \in (1, 2)$;

- $\Phi_G(Q)$ has a zero of multiplicity 1 at $Q = 2$.
- $\Phi_G(Q)$ is non-zero with sign $(-1)^{|E|-|V|+1}$ for $Q \in (2, \delta)$, where $\delta \approx 2.546$ is the flow root of the cube in the interval $(2, 3)$ [i.e., the zero in this interval of $Q^3 - 9Q^2 + 29Q - 32$].

The generalised Petersen graphs $G(nk, k)$ satisfy $|V| = 2nk$, $|E| = 3nk$, and $b = 1$. Therefore, the above theorems imply that, for any $n > 2$, $\Phi_G(Q)$ has only two simple real roots in the interval $(-\infty, \delta)$: $Q = 1, 2$. The sign for $Q \in (-\infty, 1) \cup (2, \delta)$ is that of $(-1)^{nk+1}$, and it has the opposite sign for $Q \in (1, 2)$. For large enough $Q > 0$, the sign of $\Phi_G(Q)$ is always positive. Therefore, for even k and any $n \geq 3$, or for odd k and even $n \geq 4$, this implies the existence of a real zero in $[\delta, \infty)$.⁸

We have explicitly checked that all the computed flow polynomials $\Phi_{G(nk, k)}$ have only two simple roots in the whole interval $(-\infty, \delta)$, namely $Q = 1, 2$. Furthermore, for all even (resp. odd) k , and all $n \geq 3$ (resp. all even $n \geq 4$), the polynomial $\Phi_{G(nk, k)}$ has at least one root in $[\delta, \infty)$.

If G is a cubic graph, one can easily see whether $Q = 3$ is a flow root or not [17, Proposition 6.4.2]:

Theorem 4.9 *A bridgeless cubic graph G has a nowhere zero 3-flow if and only if it is bipartite.*

As the graphs $G(nk, k)$ are bipartite if and only if k is odd and n is even, Theorem 4.9 implies that $\Phi_{G(nk, k)}(Q)$ has at least one factor $Q - 3$ whenever $G(nk, k)$ is not bipartite: namely, when k is even or n is odd (or both). When $G(nk, k)$ is bipartite, then $\Phi_{G(nk, k)}(3) > 0$. We have checked these facts in all the flow polynomials we have explicitly computed in this work.

5 The Beraha–Kahane–Weiss theorem

A central role in the subsequent analysis (see sections 6–7) is played by a theorem on analytic functions due to Beraha, Kahane and Weiss (BKW) [8–11] and generalised slightly by Sokal [48]. The situation is as follows: Let D be a domain (connected open set) in the complex plane, and let $\alpha_1, \dots, \alpha_M, \mu_1, \dots, \mu_M$ ($M \geq 2$) be analytic functions on D , none of which is identically zero. For each integer $n \geq 0$, define

$$f_n(z) = \sum_{k=1}^M \alpha_k(z) \mu_k(z)^n. \quad (5.1)$$

We are interested in the zero sets

$$\mathcal{Z}(f_n) = \{z \in D: f_n(z) = 0\} \quad (5.2)$$

⁸The lower bound is sharp, as the cube is isomorphic to the generalised Petersen graph $G(4, 3)$, which has a zero at $Q = \delta$.

and in particular in their limit sets as $n \rightarrow \infty$:

$$\liminf \mathcal{Z}(f_n) = \{z \in D: \text{every neighbourhood } U \ni z \text{ has a nonempty intersection with all but finitely many of the sets } \mathcal{Z}(f_n)\} \quad (5.3)$$

$$\limsup \mathcal{Z}(f_n) = \{z \in D: \text{every neighbourhood } U \ni z \text{ has a nonempty intersection with infinitely many of the sets } \mathcal{Z}(f_n)\} \quad (5.4)$$

Let us call an index k *dominant at z* if $|\mu_k(z)| \geq |\mu_l(z)|$ for all l ($1 \leq l \leq M$); and let us write

$$D_k = \{z \in D: k \text{ is dominant at } z\}. \quad (5.5)$$

Then the limiting zero sets can be completely characterised as follows:

Theorem 5.1 (Beraha–Kahane–Weiss [8–11, 48]) *Let D be a domain in $\mathbb{C}$, and let $\alpha_1, \dots, \alpha_M, \mu_1, \dots, \mu_M$ ($M \geq 2$) be analytic functions on D , none of which is identically zero. Let us further assume a “no-degenerate-dominance” condition: there do not exist indices $k \neq k'$ such that $\mu_k \equiv \omega \mu_{k'}$ for some constant ω with $|\omega| = 1$ and such that $D_k (= D_{k'})$ has nonempty interior. For each integer $n \geq 0$, define f_n by*

$$f_n(z) = \sum_{k=1}^M \alpha_k(z) \mu_k(z)^n.$$

Then $\liminf \mathcal{Z}(f_n) = \limsup \mathcal{Z}(f_n)$, and a point z lies in this set if and only if either

- (a) *There is a unique dominant index k at z , and $\alpha_k(z) = 0$; or*
- (b) *There are two or more dominant indices at z .*

Note that case (a) consists of isolated points in D , while case (b) consists of curves (plus possibly isolated points where all the μ_k vanish simultaneously). Henceforth we shall denote by $\mathcal{B}$ the locus of points satisfying condition (b).

We shall often refer to the functions μ_k as “eigenvalues”, and to the α_k as “amplitudes”, because that is exactly how they arise in the transfer matrix formalism.

In Ref. [11, p. 55], Beraha, Kahane, and Weiss give (without proof) the following corollary based on their proof of Theorem 5.1:

Corollary 5.2 *If z is a real isolated limiting point, the polynomials f_n have real coefficients, the dominant eigenvalue $\mu_\star$ and its coefficient $\alpha_\star$ are both real in an interval $(z - \epsilon, z + \epsilon)$ for some $\epsilon > 0$, and if $\alpha'_\star(z) \neq 0$, then z is the limit of a real sequence $\{z_n\}$ with $f_n(z_n) = 0$.*

The proof is simple: For any sufficiently small $\epsilon > 0$, $\alpha_\star(z \pm \epsilon)$ are non-zero of opposite sign, and still dominant. Then, for all sufficiently large n (depending on ϵ), $f_n(z \pm \epsilon)$ are real of opposite sign. Therefore there exists a root in between.

In the next section we need some general results on the existence of a real sequence of zeros $\{z_n\}$ such that it converges to a real non-isolated limiting point. These results can be summarised in the following

Lemma 5.3 *Let us suppose that z is a real non-isolated limiting point, such that exactly two dominant eigenvalues μ_1 and μ_2 become equimodular at z . Let us further suppose that:*

- (a) *The two dominant eigenvalues are analytic functions in a neighbourhood of z .*
- (b) *The eigenvalue μ_1 (resp. μ_2) is dominant (resp. subdominant) in the interval $[z - \epsilon, z)$, and is subdominant (resp. dominant) in the interval $(z, z + \epsilon]$ for some $\epsilon > 0$.*
- (c) *The corresponding amplitudes α_1 and α_2 do not vanish at z .*
- (d) *The eigenvalues μ_1, μ_2 and the amplitudes α_1, α_2 are real in a real neighbourhood of z .*

Then,

- 1. *If $\alpha_1(z)\alpha_2(z) > 0$ and $\mu_1(z) = -\mu_2(z)$, then for odd n , there is a real sequence of zeros $\{z_n\}$ converging to z .*
- 2. *If $\alpha_1(z)\alpha_2(z) < 0$ and $\mu_1(z) = -\mu_2(z)$, then for even n , there is a real sequence of zeros $\{z_n\}$ converging to z .*
- 3. *If $\alpha_1(z)\alpha_2(z) < 0$ and $\mu_1(z) = \mu_2(z)$, then for all n , there is a real sequence of zeros $\{z_n\}$ converging to z .*

PROOF. The function f_n (5.1) can be written as

$$f_n = \alpha_1 \mu_1^n + \alpha_2 \mu_2^n + \sum_{j=3}^N \alpha_j \mu_j^n, \quad (5.6)$$

where the $N - 2$ other eigenvalues are subdominant in a neighbourhood of z . Then, for every sufficiently small $\epsilon > 0$ we can find real numbers $0 < r_1, r_2 < 1$ such that

$$\left| \frac{\mu_i(w)}{\mu_1(w)} \right| \leq r_1, \quad (5.7)$$

for all $i \geq 2$ and $w \in [z - \epsilon, z)$, and

$$\left| \frac{\mu_i(w)}{\mu_2(w)} \right| \leq r_2, \quad (5.8)$$

for all $i = 1$ or $i \geq 3$ and $w \in (z, z + \epsilon]$. Let M the maximum of the amplitudes α_i in the interval $[z - \epsilon, z + \epsilon]$. We choose ϵ small enough so that the signs of the dominant eigenvalues and amplitudes is the same as at z . Then, we have that

$$f_n(z - \epsilon) = \mu_1^n \left[\alpha_1 + \sum_{j=2}^N \alpha_j \left(\frac{\mu_j}{\mu_1} \right)^n \right] \quad (5.9a)$$

$$f_n(z + \epsilon) = \mu_2^n \left[\alpha_2 + \alpha_1 \left(\frac{\mu_1}{\mu_2} \right)^n + \sum_{j=3}^N \alpha_j \left(\frac{\mu_j}{\mu_2} \right)^n \right] \quad (5.9b)$$

Then, for large enough n , the quantities $f_n(z \pm \epsilon)$ have opposite sign in the following cases:

1. $\alpha_1\alpha_2 > 0$ and $(\mu_1/\mu_2)^n < 0$, which occurs if $\mu_1\mu_2 < 0$ and n is odd.
2. $\alpha_1\alpha_2 < 0$ and $(\mu_1/\mu_2)^n > 0$, which occurs if $\mu_1\mu_2 > 0$ (then n can have either parity) or if $\mu_1\mu_2 < 0$ and n is even.

In any of these cases, the continuous function f_n attains values of distinct signs at the endpoints of the interval $[z - \epsilon, z + \epsilon]$, therefore there should be a zero at some point inside this interval. ■

Remarks. 1. In the fourth case $\alpha_1\alpha_2 > 0$ and $\mu_1 = \mu_2$, the zeros converging to z are *non-real*.

2. If the derivative of the ratio μ_1/μ_2 is nonvanishing at z , then condition (b) [or the same condition with μ_1 and μ_2 interchanged] necessarily holds. However, the converse is false: it is possible for condition (b) to hold even if μ_1/μ_2 has a vanishing derivative at z .

As remarked on Ref. [11, p. 55], the convergence ratio for isolated and non-isolated limiting points is rather different: exponentially fast for the former $|z_n - z| \leq Ar^n$, and $|z - z_n| \leq An^{-1}$ for the latter, as $n \rightarrow \infty$.

6 Real zeros of the flow polynomials $\Phi_{G(nk,k)}$

In this section we will discuss the real zeros of the flow polynomials $\Phi_{G(nk,k)}$ for $1 \leq k \leq 7$. In particular, we will focus on the real zeros around $Q = 5$, and on the existence of real zeros $Q > 5$.

6.1 $k \leq 5$

For $1 \leq k \leq 3$, we compute the flow polynomials of $G(nk, k)$ and their roots for all n in the range $1 \leq n \leq 30$. All the flow roots we have found are smaller than $Q = 4$. We conjecture that this holds for larger n as well. For $k = 4, 5$ we find flow polynomials with real roots greater than $Q = 4$:

- $G(28, 4)$ has two real roots greater than $Q = 4$: $Q_1 \approx 4.0002086861$ and $Q_2 \approx 4.3876416603$. As n grows, the maximal real root of $\Phi_{G(4n,4)}$ tends to the value $Q_c(4) \approx 4.5697435537$.
- $G(30, 5)$ has two real roots greater than $Q = 4$: $Q_1 \approx 4.0000786673$ and $Q_2 \approx 4.4867394006$. As n grows, the maximal real root of $\Phi_{G(5n,5)}$ tends to the value $Q_c(5) \approx 4.9029018077$.

Both families thus provide counter-examples to Welsh's conjecture (cf. Conjecture 1.3).

Remark. The values of $Q_c(k)$ mentioned above are obtained by determining the eigenvalue crossing that corresponds to case (b) of Theorem 5.1. See Conjecture 7.2 below for a more precise statement about the leading eigenvalues at $Q_c(k)$.

6.2 $k = 6, 7$: The isolated limiting point $Q = 5$

The family $G(6n, 6)$ also provides a counter-example to Conjecture 1.3: it displays a real zero converging to $Q = 5$ from below; for instance, $G(144, 6)$ has a real zero at $Q \approx 4.9987003379$.

This empirical observation will be made rigorous by applying Corollary 5.2 to the family $G(6n, 6)$ at $Q = 5$. In this case, there is a unique (and therefore, real) leading eigenvalue, which corresponds to $\ell = 3$ and $\lambda = (3)$: $\mu_{3,(3),*}(5) \approx 177.122$. The corresponding amplitude $\alpha_{3,(3)}$ is given by (3.14),

$$\alpha_{3,(3)}(Q) = \frac{1}{8}Q(Q-1)(Q-5), \quad (6.1)$$

and is a polynomial in Q with a single zero at $Q = 5$. Therefore, the BKW theorem implies that $Q = 5$ is an isolated limiting point for this family. The fact that the limiting point $Q = 5$ is isolated implies that there is a interval of radius ϵ around $Q = 5$ where there are no other limiting points. Therefore, in this interval the eigenvalue $\mu_{3,(3),*}$ is still dominant, and hence real. Therefore, Corollary 5.2 implies that there is a sequence of *real* zeros $\{Q_n\}$ converging to $Q = 5$. We can go a little bit further and show that:

Corollary 6.1 *The point $Q = 5$ is an isolated limiting point for the family $G(6n, 6)$ with $n \geq 3$. There is a sequence of real zeros $\{Q_n\}$ of the flow polynomial $\Phi_{G(6n,6)}$ that converges to $Q = 5$ from below.*

PROOF. Let us consider the point $Q = 5 - \epsilon$ with ϵ small enough so that: 1) the leading eigenvalue $\mu_{3,(3),*}(Q) > 0$; 2) all other N sub-leading eigenvalues μ_j satisfy $|\mu_j(Q)/\mu_{3,(3),*}(Q)| \leq r < 1$; 3) the sub-leading amplitudes $|\alpha_j| \leq M$ are bounded; and 4) $\alpha_{3,(3)}(Q) < 0$. We can always choose ϵ so that these conditions are fulfilled, as the eigenvalues and amplitudes are analytic functions of Q in a neighbourhood of $Q = 5$. In addition, we know that $\Phi_{G(6n,6)}(5) > 0$, due to Lemma 3.1. Therefore, we only need to show that $\Phi_{G(6n,6)}(Q) < 0$ for n large enough. This is easy as,

$$\Phi_{G(6n,6)}(Q) = \mu_{3,(3),*}(Q)^n \left[\alpha_{3,(3)}(Q) + \sum_{j=1}^N \alpha_j(Q) \left(\frac{\mu_j(Q)}{\mu_{3,(3),*}(Q)} \right)^n \right]. \quad (6.2)$$

We can always choose N_0 such that $NMr^{N_0} < |\alpha_{3,(3)}(5 - \epsilon)|$. Then, for all $n \geq N_0$, the sign of $\Phi_{G(6n,6)}(Q)$ is that of $\alpha_{3,(3)}(Q)$ (i.e., negative), so that there should be a point Q_0 in $(5 - \epsilon, 5)$ such that $\Phi_{G(6n,6)}(Q_0) = 0$. ■

The same situation applies to the family $G(7n, 7)$ at $Q = 5$. In this case, the unique (and real) leading eigenvalue is $\mu_{3,(3),*}(5) \approx -621.779$. Therefore, $Q = 5$ is an isolated limiting point for this family, and there is a *real* sequence of zeros $\{Q_n\}$ converging to $Q = 5$. Because $\mu_{3,(3),*}(5) < 0$, the above arguments imply that the convergence of the sequence $\{Q_n\}$ to $Q = 5$ is a bit more complicated:

Corollary 6.2 *The point $Q = 5$ is an isolated limiting point for the family $G(7n, 7)$ with $n \geq 3$. There is a sequence of real zeros $\{Q_n\}$ of the flow polynomial $\Phi_{G(7n,7)}$ converging to $Q = 5$. The sub-sequence with odd n (resp. even n) converges to $Q = 5$ from above (resp. below).*

PROOF. If we consider even n , then the proof is as before; therefore the sequence $\{Q_{2p}\}_{p \in \mathbb{N}}$ converges to $Q = 5$ from below. On the contrary, for odd n , then the sign of the leading term $\alpha_{3,(3)}\mu_{3,(3),*}(Q)^n$ is positive for $Q < 5$, and negative for $Q > 5$. Therefore, a trivial modification of the above arguments leads to the convergence of the sub-sequence $\{Q_{2p+1}\}_{p \in \mathbb{N}}$ to $Q = 5$ from above. ■

Indeed, Corollaries 6.1 and 6.2 imply Theorem 1.7. Moreover, Corollary 6.2 implies the existence of real flow roots arbitrarily close to $Q = 5$ on *both* sides (above and below).

6.3 $k = 7$: Real flow-polynomial zeros larger than $Q = 5$

There are also real non-isolated limiting points for the families $G(nk, k)$ with $1 \leq k \leq 7$. These non-isolated limiting points corresponds to “crossings” between two dominant eigenvalues, each of them coming from a block of the transfer matrix $\widehat{T}_{k+1,\ell,\lambda}$ with different values of ℓ . Therefore, both dominant eigenvalues are real and analytic in some real interval around the non-isolated limiting point $Q_c(k)$. In some cases, there corresponds a sequence of real zeros $\{Q_n\}$ converging to that limiting point.

The family $G(6n, 6)$ would be in principle a good candidate for having real roots larger than $Q = 5$: a direct calculation shows that $Q_c(6) \approx 5.1079785012$ is a non-isolated limiting point for this family. However, the actual computation of all the members of this family up to $G(144, 6)$ does not reveal any zero $Q > 5$.

The explanation is simple: at $Q = Q_c(6)$ both dominant eigenvalues are equal $\lambda_{1,(1),*}(Q_c(6)) = \lambda_{3,(3),*}(Q_c(6)) \approx 169.757$, and the corresponding amplitudes are both positive. Therefore, the hypotheses of Lemma 5.3 are not satisfied, and we cannot find *real* zeros converging to the non-isolated limiting point $Q_c(6)$. They should be non-real.

Finally, we have found that the family $G(7n, 7)$ with *odd* n has members with the desired property. Even though $G(63, 7)$ does not have any zeros larger than $Q = 5$, the next member that we have computed $G(119, 7)$ has two zeros: $Q_1 \approx 5.0000197675$, and $Q_2 \approx 5.1653424423$. The flow polynomial for $G(119, 7)$ has degree 120 and can be written as

$$\Phi_{G(119,7)}(Q) = (Q - 1)(Q - 2)(Q - 3)P_{117}(Q), \quad (6.3)$$

where $P_{117}(Q) = Q^{117} - 351Q^{116} + 61191Q^{115} - 7064107Q^{114} + \dots$ is a polynomial in Q of degree 117. The coefficients of this polynomial are given in Table 2. We can formalise the existence of such real roots greater than $Q = 5$ in the following way:

Proposition 6.3 *Let $\Phi_G(Q)$ be the flow polynomial of the generalised Petersen graph $G = G(119, 7)$. Then it has a real zero in the interval $(5 + 10^{-5}, 5 + 2 \times 10^{-5})$, and another real zero in the interval $(516534 \times 10^{-5}, 516535 \times 10^{-5})$.*

PROOF. We can evaluate the polynomial $\Phi_G(Q)$ at the two end-points of the interval $(5 + 10^{-5}, 5 + 2 \times 10^{-5})$ using exact rational arithmetic and find results of distinct sign: $\Phi_G(5 + 10^{-5}) \approx +2.21791 \times 10^{42}$, and $\Phi_G(5 + 2 \times 10^{-5}) \approx -5.27937 \times 10^{40}$. Therefore, the intermediate value theorem ensures the existence of a zero of Φ_G in the open interval $(5 + 10^{-5}, 5 + 2 \times 10^{-5})$.

The same procedure can be carried out for the second interval: $\Phi_G(516534 \times 10^{-5}) \approx -1.46592 \times 10^{42}$, and $\Phi_G(516535 \times 10^{-5}) \approx +4.53729 \times 10^{42}$. ■

Remark. As a curiosity, $\Phi_{G(119,7)}(5) = 4488918995790513676672232799446257724715600$, and $\Phi_{G(119,7)}(4) = 1133172760943853528$.

As a matter of fact, we also expect that the family $G(7n, 7)$ has, for every large enough odd n , two real zeros larger than $Q = 5$: one converging to $Q = 5$ from above (as we have discussed in the previous section), and the other one converging to the limiting point $Q_c(7) \approx 5.2352605291$ from below. The complete statement is given by the following

Lemma 6.4 *The point $Q_c(7) \in (5235260 \times 10^{-6}, 5235261 \times 10^{-6})$ is a non-isolated limiting point for the family $G(7n, 7)$ with $n \geq 3$. There is a sequence of real zeros $\{Q_n\}$ of the flow polynomial $\Phi_{G(7n,7)}$ converging to $Q_c(7)$ from below for odd n .*

PROOF. In the interval $[5235260 \times 10^{-6}, 5235261 \times 10^{-6}]$, we only find two dominant eigenvalues $\lambda_{0,*}$ and $\lambda_{3,(3),*}$, each of them coming from a different ℓ sector. Therefore, both eigenvalues and its sum $\lambda_{0,*} + \lambda_{3,(3),*}$ are analytic functions of Q in this interval. If we evaluate $\lambda_{0,*} + \lambda_{3,(3),*}$ at the two end-points of this interval using MATHEMATICA, we find results of distinct sign: $\lambda_{0,*}(5235260 \times 10^{-6}) + \lambda_{3,(3),*}(5235260 \times 10^{-6}) \approx 0.000917$, and $\lambda_{0,*}(5235261 \times 10^{-6}) + \lambda_{3,(3),*}(5235261 \times 10^{-6}) \approx -0.000817$. Therefore, there exists an intermediate value $Q_c(7)$ in the open interval $(5235260 \times 10^{-6}, 5235261 \times 10^{-6})$ such that $\lambda_{0,*}(Q_c(7)) + \lambda_{3,(3),*}(Q_c(7)) = 0$. Exactly at $Q = Q_c(7)$, these two eigenvalues have opposite signs $\lambda_{0,*}(Q_c(7)) = -\lambda_{3,(3),*}(Q_c(7)) \approx -565.833$. The corresponding amplitudes are both positive for any $Q > 5$. Therefore, according to Lemma 5.3 we find real zeros converging to $Q_c(7)$ only for large enough odd values of n .

The fact that the convergence is from below comes from the fact that $\alpha_{0,*}(Q_c(7)) - \alpha_{3,(3),*}(Q_c(7)) > 0$, so the sign of the flow polynomial at $Q_c(7)$ is positive $\Phi_{G(7n,7)}(Q_c(7)) \approx [\alpha_{0,*}(Q_c(7)) - \alpha_{3,(3),*}(Q_c(7))] \lambda_{0,*}(Q_c(7))^n > 0$. However, its sign at $Q_c(7) - \epsilon$ (for small enough values of $\epsilon > 0$) $\Phi_{G(7n,7)}(Q_c(7) - \epsilon) \approx \alpha_{3,(3),*}(Q_c(7)) \lambda_{3,(3),*}(Q_c(7))^n < 0$ for odd n . Therefore, there should be a root in between. ■

Proposition 6.3 and Lemma 6.4 imply Theorem 1.8.

7 Numerical results

We now turn to the computation of the limiting curves $\mathcal{B}_k$. For any fixed integer $k \geq 1$, the curve $\mathcal{B}_k$ is defined as the set of non-isolated limiting points of the zeros of the flow polynomial $\Phi_{G(nk,k)}$, as $n \rightarrow \infty$. We have computed these curves for $k = 1, 2, \dots, 7$ using the direct-search method, as described in [26]. The results are shown in Figures 2–5, both in the complex Q -plane and in the complex $1/Q$ -plane.

We have noticed several empirical patterns in Figures 2–5. The first one is related to the number of outward branches of the limiting curve $\mathcal{B}_k$, and the dominant eigenvalue in each asymptotic sector:⁹

⁹This is exactly the same empirical behavior found for the limiting curves $\mathcal{B}_m$ associated to the chromatic zeros of the family $S_{m,n}$ when $n \rightarrow \infty$ [43, Conjeture 7.1]. The graph $S_{m,n}$ can be regarded as a square-lattice grid with m columns, n rows, free boundary conditions in the longitudinal direction, and special boundary conditions in the transverse direction: we introduce two extra vertices such that every vertex on the leftmost column of the grid is connected to one of them, and every vertex on the rightmost column is connected to the other one.

Conjecture 7.1 Fix $k \geq 1$. Then, the limiting curve $\mathcal{B}_k$ has exactly $2k$ outward branches extending to $Q = \infty$, with asymptotic angles $\arg Q = \theta_{k,j}$ where

$$\theta_{k,j} = \left(m - \frac{1}{2}\right) \frac{\pi}{k}, \quad \text{for } j = 1, \dots, 2k. \quad (7.1)$$

Moreover, the dominant eigenvalue is $\mu_{k+1,0,\star}$ in the asymptotic regions

$$\theta_{k,j} \leq \arg Q \leq \theta_{k,j+1} \quad \text{for} \quad \begin{cases} j = 1, 3, \dots, 2k-1 & \text{if } k \text{ is even} \\ j = 2, 4, \dots, 2k & \text{if } k \text{ is odd} \end{cases} \quad (7.2)$$

while in the other asymptotic sectors the dominant eigenvalue is $\mu_{k+1,1,(1),\star}$.

We may conjecture that, as $k \rightarrow \infty$, the limiting curves $\mathcal{B}_k$ (without the outward branches) converge to some curve $\mathcal{B}_\infty$. This is what Figure 6 suggests. Indeed, outside this curve, the outward branches seem to get denser as k increases. Therefore, we conjecture that the set of accumulation points of the flow-polynomial zeros for the generalised Petersen graphs $G(nk, k)$ is *dense* in the whole complex Q -plane, except in the interior of the curve $\mathcal{B}_\infty$. If this conjecture is true, it implies in particular that flow roots of cubic graphs are dense in the complex plane, except possibly in some bounded region near the origin. Note that a similar result holds (by duality) for the flow roots of planar graphs [48].

If we restrict to the cases with $k \geq 3$ (see Figures 3–5), we notice some regularities about the isolated limiting points and the dominant eigenvalues in the non-asymptotic region of the complex q -plane:

Conjecture 7.2 Fix $k \geq 4$. Then, the dominant eigenvalue in the regions that intersect the real Q -axis is

$$\mu_\star = \begin{cases} \mu_{k+1,1,(1),\star} & \text{for } \operatorname{Re} Q \in (-\infty, 2] \\ \mu_{k+1,2,(2),\star} & \text{for } \operatorname{Re} Q \in [2, 4] \\ \mu_{k+1,3,(3),\star} & \text{for } \operatorname{Re} Q \in [4, Q_c(k)] \\ \mu_{k+1,1,(1),\star} & \text{for } \operatorname{Re} Q \in [Q_c(k), \infty) \text{ and even } k \\ \mu_{k+1,0,\star} & \text{for } \operatorname{Re} Q \in [Q_c(k), \infty) \text{ and odd } k \end{cases} \quad (7.3)$$

Therefore, $Q = 1$, $Q = 3$ and $Q = 5$ (only when $Q_c(k) > 5$) are isolated limiting points; and $Q = 2$, $Q = 4$, and $Q = Q_c(k)$ are non-isolated limiting points.

Notice that the above conjecture is also valid for $k = 3$, except that in this case, there is no region where the eigenvalue $\mu_{k+1,3,(3),\star}$ becomes dominant, as $Q_c(3) < 4$. It is worth noting that the dominant eigenvalue comes from the completely symmetric irreducible representation of S_ℓ for $\ell = 1, 2, 3$.

Assuming that the dominant eigenvalue on the real Q -axis invariably comes from the completely symmetric representation, it is possible to extend the numerical determination of $Q_c(k)$ to higher values of k . The relevant transfer matrices have dimension $(\ell!)^{-1} |A_{k+1}^{(\ell)}|$ and can be diagonalised by a standard iterative scheme that uses their decomposition as a product of sparse matrices, as in (3.8). The resulting values of $Q_c(k)$ are shown in Table 1.

k	$Q_c(k)$
1	3
2	3.6180339887
3	3.7818423129
4	4.5697435537
5	4.9029018077
6	5.1079785012
7	5.2352605291
8	5.3246966903
9	5.3886186958
10	5.4364766073
11	5.4729804532

Table 1: Values of $Q_c(k)$ for the generalised Petersen graphs $G(nk, k)$ for $1 \leq k \leq 11$.

The values of $Q_c(k)$ are seen to be well fitted by a second-degree polynomial in $1/k$, provided that data for k even and k odd are fitted separately. Both fits are consistent with each other and give the common limit

$$Q_c \equiv \lim_{k \rightarrow \infty} Q_c(k) = 5.75 \pm 0.01. \quad (7.4)$$

Formalising slightly these experimental results, we conjecture that there exists a real number $Q_c \approx 5.75$ that is an accumulation point of real flow zeros for the graph family $G(nk, k)$, in the limit $\lim_{n \rightarrow \infty} \lim_{k \rightarrow \infty}$, and that no generalised Petersen graph has a real flow zero above Q_c . The hope that no graph family fares better than the generalised Petersen graph is our main motivation—and evidence—for stating Conjecture 1.9.




Figure 2: Zeros of the flow polynomial and limiting curves $\mathcal{B}_k$ for the generalised Petersen graphs $G(nk, k)$ with $k = 1$ (a,b), and $k = 2$ (c,d). In (a,c) we show these results in the complex Q -plane, while in (b,d), we use the complex $1/Q$ -plane. For each k , we show the flow-polynomial zeros of the generalised Petersen graphs $G(12k, k)$ ($\square$ black), $G(16k, k)$ ($\circ$ red), $G(20k, k)$ ($\triangle$ green), and $G(24k, k)$ ($\diamond$ blue).




Figure 3: Zeros of the flow polynomial and limiting curves $\mathcal{B}_k$ for the generalised Petersen graphs $P(nk, k)$ with $k = 3$ (a,b), and $k = 4$ (c,d). In (a,c) we show these results in the complex Q -plane, while in (b,d), we use the complex $1/Q$ -plane. For each k , we show the flow-polynomial zeros of the generalised Petersen graphs $P(12k, k)$ ($\square$ black), $P(16k, k)$ ($\circ$ red), $P(20k, k)$ ($\triangle$ green), and $P(24k, k)$ ($\diamond$ blue).




Figure 4: Zeros of the flow polynomial and limiting curves $\mathcal{B}_k$ for the generalised Petersen graphs $P(nk, k)$ with $k = 5$ (a,b), and $k = 6$ (c,d). In (a,c) we show these results in the complex Q -plane, while in (b,d), we use the complex $1/Q$ -plane. For each k , we show the flow-polynomial zeros of the generalised Petersen graphs $P(12k, k)$ ($\square$ black), $P(16k, k)$ ($\circ$ red), $P(20k, k)$ ($\triangle$ green), and $P(24k, k)$ ($\diamond$ blue).




Figure 5: Zeros of the flow polynomial and limiting curves $\mathcal{B}_7$ for the generalised Petersen graphs $P(7n, 7)$. In (a) we show these results in the complex Q -plane, while in (b), we use the complex $1/Q$ -plane. We show the flow-polynomial zeros of the generalised Petersen graphs $P(21, 7)$ ($\square$ black), $P(35, 7)$ ($\circ$ red), $P(63, 7)$ ($\triangle$ green), and $P(119, 7)$ ($\diamond$ blue).

8 Discussion

In this concluding section we would like to place our results on the flow polynomial in a larger perspective, and to propose a number of conjectures that should stimulate further research. We have seen that the Q -state Potts model with temperature parameter v comprises both the flow polynomial ($v = -Q$) and the chromatic polynomial ($v = -1$) as special cases. It is therefore useful to compare our results on the flow polynomial for the particular class of non-planar graphs $G(n, k)$ with what is known for the chromatic polynomial for certain classes of planar graphs. More generally, we should like to outline and compare the results and expectations for real accumulation points of chromatic zeros for planar graphs, and of flow zeros for non-planar graphs, when the graphs are taken to consist of n identical layers of width L vertices and we impose periodic boundary conditions in the n -direction. Going beyond this, we discuss also the behaviour of real accumulation points of zeros for the Potts-model partition function $Z_G(Q, v)$ along more general curves in the (Q, v) plane.

Since many of the recent results on chromatic zeros have been obtained at the borderline between mathematics and theoretical physics, we shall use—for lack of a better term—the denomination “result” to characterise a fact which has been derived using exact (but non-rigorous in the mathematical sense) methods of theoretical physics. These methods include in particular two-dimensional conformal field theory and integrability techniques, and renormalisation group arguments.

Our main objective is to spell out a number of striking analogies between, on the one hand, the intersection of the limiting curves of Figure 6 with the real Q -axis, and Conjecture 7.2 on the dominance pattern of eigenvalues on the real Q -axis, and, on the other hand, the corresponding results [26] for the chromatic zeros for cyclic strips of regular planar lattices.




Figure 6: Limiting curves $\mathcal{B}_k$ for the generalised Petersen graphs $P(kn, k)$ in the limit $n \rightarrow \infty$. In (a) we show these results in the complex Q -plane, while in (b), we use the complex $1/Q$ -plane. We show the curves corresponding to $k = 1$ (black), $k = 2$ (red), $k = 3$ (green), $k = 4$ (blue), $k = 5$ (magenta), $k = 6$ (orange), and $k = 7$ (gray).

In the planar chromatic case the general situation is expected to be the following:¹⁰

Conjecture 8.1 *Consider a family $G_{n,L}$ of planar connected graphs, consisting of n identical layers of width L vertices, with periodic boundary conditions in the n -direction. Let $\mathcal{A}_L$ (resp. $\mathcal{B}_L$) denote the set of limiting points of chromatic zeros for $G_{n,L}$, in the limit $n \rightarrow \infty$, that satisfies condition (a) (resp. condition (b)) of Theorem 5.1. Further let B_p be the p -th Beraha number:*

$$B_p = 4 \cos^2 \left(\frac{\pi}{p} \right), \quad p \in \mathbb{N}, \quad (8.1)$$

and let

$$B^{\text{even}} = \{Q = B_p : p \in 2\mathbb{N}\}, \quad B^{\text{odd}} = \{Q = B_p : p \in 2\mathbb{N} + 1\}$$

denote, respectively, the set of even and odd Beraha numbers B_p . Then:

1. There exists, for each L , a number $Q_c(L) \in (0, 4)$ so that

$$\begin{aligned} (a) \quad \mathcal{A}_L \cap \mathbb{R} &= B^{\text{odd}} \cap [0, Q_c(L)) \\ (b) \quad \mathcal{B}_L \cap \mathbb{R} &= B^{\text{even}} \cap [0, Q_c(L)) \cup \{Q_c(L)\} \end{aligned}$$

2. For any $p \in \mathbb{N}$, the dominant eigenvalue of the transfer matrix for

$$Q \in (B_{2p}, B_{2(p+1)}) \cap [0, Q_c(L))$$

comes from the sector with p marked clusters.

¹⁰In this section, B_p denotes the Beraha numbers (8.1), not the Bell numbers B_n (3.10) defined earlier.

Compelling evidence for Conjecture 8.1 comes from the

Proposition 8.2 (Jacobsen-Salas [26]) *Conjecture 8.1 is correct for cyclic strips of the square and triangular lattices with $L \leq 8$. The corresponding values of $Q_c(L)$ can be found in [26].*

From the point of view of statistical physics, the natural step suggested by these finite- L results is to go to the so-called infinite-volume limit, $\lim_{L \rightarrow \infty}$. We have then the following non-rigorous “results” obtained by the methods of theoretical physics:

Result 8.3 [26, 44, 45] *For each family of planar graphs $G_{n,L}$, defined as above, the limit $Q_c = \lim_{L \rightarrow \infty} Q_c(L)$ exists and satisfies $0 < Q_c \leq 4$.*

Result 8.4 [4–6, 28, 29] *For cyclic strips of the square lattice, $Q_c = 3$. For cyclic strips of the triangular lattice, $Q_c = 2 + \sqrt{3}$.*

The deep meaning of these results is that the special role of the Beraha numbers in the production of real chromatic zeros is brought out whenever the chromatic line ($v = -1$) intersects the so-called Berker-Kadanoff (BK) phase in the (Q, v) plane of the Potts model [44, 45]. For each graph family, and for each $Q \in [0, 4)$, it can be argued [30] that the extent of the BK phase is a finite non-empty interval $v_-(Q) < v < v_+(Q)$ with the properties $v_+(0) = 0$ and $\lim_{Q \rightarrow 4} v_+(Q) = \lim_{Q \rightarrow 4} v_-(Q)$. In particular, other choices of the temperature curve $v(Q)$ are possible and might in some cases lead to higher values of Q_c than those in Result 8.4. The bound on Q_c given in Result 8.3 is even sharp:

Result 8.5 *There exists a family of planar graphs $G_{n,L}$, defined as above, and a curve $v = f(Q)$ in the (Q, v) plane, so that the limiting set of zeros for the Potts-model partition function $Z_G(Q, v)$ satisfies Conjecture 8.1 and Result 8.3 with $Q_c = 4$.*

Indeed, this is the case [30, 44, 45] for cyclic strips of the square lattice with the choice [2] $v^2 = Q$ for $0 \leq Q \leq 4$ and $-2 \leq v \leq 0$. Another example is provided by cyclic strips of the triangular lattice with the choice [3] $v^3 + 3v^2 = Q$ for $0 \leq Q \leq 4$ and $-2 \leq v \leq 0$.

Let us finally mention that Conjecture 8.1 enjoys strong support from the special representation theory of the quantum group at roots of unity [44, 45].

Turning now to the flow zeros of non-planar graphs, Figure 6 and Conjecture 7.2 makes us confident that the statements of Conjecture 8.1 hold true also in the flow case, provided that one replaces even/odd Beraha numbers by even/odd integers. On a fundamental level, the replacement of Beraha numbers by integers stands out most clearly by comparing the eigenvalue amplitudes in the two cases [see, e.g., [26, Eqs. (2.28)–(2.29)] and Eq. (3.14)]. To be precise:

Conjecture 8.6 *Consider a family $G_{n,L}$ of non-planar connected graphs, consisting of n identical layers of width L vertices, with periodic boundary conditions in the n -direction. Let $\mathcal{A}_L$ (resp. $\mathcal{B}_L$) denote the set of limiting points of flow zeros for $G_{n,L}$, in the limit $n \rightarrow \infty$, that satisfies condition (a) (resp. condition (b)) of Theorem 5.1. Then:*

1. *There exists, for each L , a number $Q_c(L) > 0$ so that*

- (a) $\mathcal{A}_L \cap \mathbb{R} = (2\mathbb{N} - 1) \cap [0, Q_c(L))$
- (b) $\mathcal{B}_L \cap \mathbb{R} = (2\mathbb{N}) \cap [0, Q_c(L)) \cup \{Q_c(L)\}$

2. For any $p \in \mathbb{N}$, the dominant eigenvalue of the transfer matrix for

$$Q \in (2(p-1), 2p) \cap [0, Q_c(L))$$

comes from the fully symmetric irreducible representation (p) of the sector with p marked clusters.

This conjecture has been validated in this paper for $G_{n,L} = G(nk, k)$, the generalised Petersen graphs, with $L = k + 1$ and $k \leq 7$. To parallel Results 8.3–8.4, we have presented numerical evidence that the limit Q_c exists in this case, with the value (7.4).

Improving on Conjecture 8.6 we can characterise more fully the structure of the eigenvalues and their crossings.

Conjecture 8.7 *Consider a family $G_{n,L}$ of non-planar connected graphs, as above. Let Q_ℓ denote the smallest $Q > 0$ so that the transfer matrix $\tilde{T}_{L,\ell,(\ell)}$ —corresponding to the fully symmetric representation (ℓ) of ℓ marked blocks—has two or more dominant eigenvalues, or let $Q_\ell = \infty$ if no such Q exists. For $Q \in (0, Q_\ell)$, let λ_ℓ denote the corresponding (unique) dominant eigenvalue. Then:*

1. $Q_0 < Q_1 < Q_2 < \dots$.
2. λ_ℓ is real.
3. The equation $|\lambda_\ell| = |\lambda_{\ell'}|$ has at most one solution for $0 < Q < \min(Q_\ell, Q_{\ell'})$.
4. If such a solution exists, it is $Q = \ell + \ell' - 1$.

This conjecture has been validated in this paper for $G_{n,L} = G(nk, k)$, with $L = k + 1$ and $k \leq 7$.

There is solid evidence [6, 28, 41] that the planar graphs with the largest possible real chromatic roots are large pieces of the triangular lattice. However, when cyclic boundary conditions are imposed, even such lattices cannot saturate the bound in Result 8.3 (i.e., have $Q_c = 4$), as we have seen in Result 8.4. Replacing now chromatic roots of planar graphs with flow roots of non-planar graphs, a comparison between Conjectures 8.1 and 8.6 reveals that the special role of $Q = 4$ in the former case is replaced by $Q = \infty$ in the latter. But although we have studied the generalised Petersen graphs because they could be expected—on the basis of sporadic evidence for small graphs [22]—to have the largest possible real flow roots, we should not forget that the flow polynomial is merely another special case of the Potts model. Therefore, by analogy, there seems no reason to believe that real flow roots can be arbitrarily large. This motivates, once more, Conjecture 1.9 that we have made in the introduction.

To recover the complete analogy with Result 8.5 we should therefore like to state one last

Conjecture 8.8 *There exists a family of non-planar graphs $G_{n,L}$, defined as above, and a curve $v = f(Q)$ in the (Q, v) plane, so that the limiting set of zeros for the Potts-model partition function $Z_G(Q, v)$ satisfies Conjecture 8.1, but with $\lim_{L \rightarrow \infty} Q_c(L) = \infty$.*

To conclude, we recall that for the Potts model on planar graphs $G_{n,L}$, when $Q = (2 \cos(\pi/p))^2$ is a Beraha number the representation theory of the underlying quantum algebra [38] leads to massive degeneracies in the spectrum of the transfer matrix. In particular, depending on p , complete sectors of eigenvalues are contained in other sectors in an inclusion-exclusion fashion, completely independent of the value of v .

It seems quite plausible that something similar should happen in the non-planar case, when Q is an integer. We have indeed observed such phenomena for the transfer matrix (3.8) of the Potts model on the graphs $G(n,k)$. Let us mention but the simplest example of our findings, which happens for $Q = 0$ and any v . The set of eigenvalues in the $\ell = 0$ sector is then fully contained in the set of $\ell = 1$ eigenvalues. Moreover, let W_ℓ denote the set of eigenvalues in the fully antisymmetric representation $(1, 1, \dots, 1)$ of the sector $\ell + 1$. We have then observed, for any $\ell \geq 1$, that $W_{\ell+1} \setminus W_{\ell+2} \subseteq W_\ell$.

A Proof of the lemmas related to the transfer matrices

In this appendix we will provide the proofs of the extensions of Lemma 4.1, Lemma 4.2, and Corollary 4.3 for all $k \geq 1$. We will also prove Lemma 4.6.

It is useful to rewrite the transfer matrix T_L (3.8) as the product of two operators $\mathsf{T}_L = \mathsf{H}\mathsf{V}$, given by

$$\mathsf{H} = \prod_{i=k}^1 \mathsf{H}_{0i} \mathsf{V}_0 = \mathsf{H}_{01} \mathsf{V}_0 \mathsf{H}_{02} \mathsf{V}_0 \mathsf{H}_{03} \cdots \mathsf{V}_0 \mathsf{H}_{0k} \mathsf{V}_0 \quad (\text{A.1a})$$

$$\mathsf{V} = \prod_{i=1}^k \mathsf{V}_i = \mathsf{V}_1 \mathsf{V}_2 \cdots \mathsf{V}_{k-1} \mathsf{V}_k \quad (\text{A.1b})$$

where $L = k + 1$, the operators V_i and H_{0i} are defined in (3.7) with $v = -Q$, and we have used the property $[\mathsf{V}_i, \mathsf{V}_j] = [\mathsf{D}_i, \mathsf{D}_j] = 0$ for all i, j .

A.1 Extension of Lemma 4.1

Lemma A.1 *Fix $k \geq 1$, and $\ell = 0, \dots, k + 1$. Then, the diagonal block $\widehat{\mathsf{T}}_{k+1,\ell}$ contains the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$ with multiplicity $\tilde{N}_{k,1}(\ell) \ell!$, where:*

$$\tilde{N}_{k,1}(\ell) = \begin{cases} 0 & \text{if } \ell = 0 \\ \frac{1}{(\ell-1)!} |\tilde{A}_k^{(\ell-1)}| = \sum_{p=0}^k \binom{k}{p} \left\{ \begin{matrix} p \\ \ell-1 \end{matrix} \right\} S_{k-p} & \text{if } 1 \leq \ell \leq k+1 \end{cases} \quad (\text{A.2})$$

and S_n is given in (3.33).

PROOF. Let us consider first the transfer matrix $\tilde{\mathsf{T}}_L = \mathsf{P}_L \mathsf{T}_L$ with T_L given by (A.1), and the projector P_L defined in (3.17). Let us fix $\ell \geq 1$ and a representation $\lambda \in S_\ell$ of dimensionality $\dim \lambda$. The set $\tilde{A}_L^{(\ell)}$ can be split into two disjoint sets:

- $\mathcal{C}_1^\ell$ is the set of all partitions of $\{0, 1, \dots, k\}$ with no un-marked singletons, ℓ marked clusters, and vertex 0 is a marked singleton.
- $\mathcal{C}_2^\ell$ is the set of all partitions of $\{0, 1, \dots, k\}$ with no un-marked singletons, ℓ marked clusters, and vertex 0 is not a singleton. In this case, the vertex 0 can be either marked or unmarked.

It is clear that $|\tilde{A}_L^{(\ell)}| = |\mathcal{C}_1^\ell| + |\mathcal{C}_2^\ell|$. Notice that we do not need to consider the case of vertex 0 being an un-marked singleton, as all partitions of this type do not belong to $\tilde{A}_L^{(\ell)}$.

In order to find out the structure of the transfer matrix $\tilde{T}_L$, we do not need to specify the representation λ ; therefore, we will omit this index (as well as the index k) to make the notation clearer.

Let us first consider a partition $\mathcal{P}_1 \in \mathcal{C}_1^\ell$ represented by the vector $e_{\mathcal{P}_1}$ in the space of partitions. Then,

$$\mathbf{V} e_{\mathcal{P}_1} = (-Q)^k e_{\mathcal{P}_1} + \sum_{\mathcal{P} \in \tilde{\mathcal{C}}_1^\ell} A_{\mathcal{P}} e_{\mathcal{P}}, \quad (\text{A.3})$$

where the sum is over the set of all partitions $\tilde{\mathcal{C}}_1^\ell$ of $\{0, 1, \dots, k\}$ with ℓ marked blocks, with vertex 0 a marked singleton, and such that there is at least one un-marked singleton. This is because $\mathbf{V}$ only contains the identity operator and detach operators D_i with $1 \leq i \leq k$.

If we consider a partition $\mathcal{P}_2 \in \mathcal{C}_2^\ell$, then

$$\mathbf{V} e_{\mathcal{P}_2} = (-Q)^k e_{\mathcal{P}_2} + \sum_{\mathcal{P} \in \tilde{\mathcal{C}}_1^\ell} A_{\mathcal{P}} e_{\mathcal{P}} + \sum_{\mathcal{P} \in \tilde{\mathcal{C}}_2^\ell} B_{\mathcal{P}} e_{\mathcal{P}}, \quad (\text{A.4})$$

where now $\tilde{\mathcal{C}}_2^\ell$ is the set of all partitions of $\{0, 1, \dots, k\}$ with ℓ marked blocks, with vertex 0 not a singleton, and such that there is at least one un-marked singleton. We omit the possible partition with 0 an un-marked singleton, as it belongs to the kernel of the first operator $\mathbf{V}_0$ in $\mathbf{H}$.

We now notice that the operator $\mathbf{H}_{0i}\mathbf{V}_0$ can be written as:

$$\mathbf{H}_{0i}\mathbf{V}_0 = -Q I + D_0 + Q^2 J_{0i} - Q J_{0i} D_0 \quad (\text{A.5})$$

where J_{0i} and D_0 are the join and detach operators, respectively. It is not hard to see that for all partitions $\mathcal{P} \in \tilde{A}_k^{(\ell)}$:

$$\mathbf{H}_{0i}\mathbf{V}_0 e_{\mathcal{P}} = -Q e_{\mathcal{P}} + \sum_{\mathcal{P}' \in \mathcal{C}_2^\ell} A_{\mathcal{P}'} e_{\mathcal{P}'}, \quad (\text{A.6})$$

where if $\mathcal{P} \in \mathcal{C}_2^\ell$, $\mathcal{P}'$ might coincide with $\mathcal{P}$. For partitions with at least one un-marked singleton $\mathcal{P} \in \tilde{\mathcal{C}}_1^\ell \cup \tilde{\mathcal{C}}_2^\ell$, we have

$$\mathbf{H}_{0i}\mathbf{V}_0 e_{\mathcal{P}} = -Q e_{\mathcal{P}} + \sum_{\mathcal{P}' \in \mathcal{C}_2^\ell} A_{\mathcal{P}'} e_{\mathcal{P}'} + \sum_{\mathcal{P}' \in \tilde{\mathcal{C}}_2^\ell} B_{\mathcal{P}'} e_{\mathcal{P}'}. \quad (\text{A.7})$$

Notice that when we apply the full operator $\mathbf{H}$, the partitions belonging to $\tilde{\mathcal{C}}_1^\ell \cup \tilde{\mathcal{C}}_2^\ell$ are eliminated by the application of the operator $\mathbf{P}_L$ (3.17).

We now put together the above observations, and conclude that if $\mathcal{P} \in \mathcal{C}_1^\ell$, then

$$\tilde{\mathbf{T}}_L \mathbf{e}_{\mathcal{P}} = (-Q)^{2k} \mathbf{e}_{\mathcal{P}} + \sum_{\mathcal{P}' \in \mathcal{C}_2^\ell} A_{\mathcal{P}'} \mathbf{e}_{\mathcal{P}'} . \quad (\text{A.8})$$

On the other hand, if $\mathcal{P} \in \mathcal{C}_2^\ell$, then

$$\tilde{\mathbf{T}}_L \mathbf{e}_{\mathcal{P}} = \sum_{\mathcal{P}' \in \mathcal{C}_2^\ell} A_{\mathcal{P}'} \mathbf{e}_{\mathcal{P}'} . \quad (\text{A.9})$$

This means that, if we order the partition states appropriately, the transfer matrix $\tilde{\mathbf{T}}_L$ has a block triangular form. This result holds for all representations $\lambda \in S_\ell$.

The block corresponding to the partitions in $\mathcal{C}_1^\ell$ is diagonal with all eigenvalues equal to Q^{2k} . In terms of the matrix $\hat{T}_L$ (3.20), this common eigenvalue takes the value $\mu_{k,k+1} = (-1)^k Q^{-2k} Q^{2k} = (-1)^k$. This proves the existence of the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$ of $\hat{T}_L$.

The dimension of this diagonal block should be equal to the number of partitions of the set $\{1, 2, \dots, k\}$ with no un-marked singletons, and $\ell - 1$ marked blocks. (Indeed, for $\ell = 0$, this dimension is identically zero.) But now, as we are considering linear combinations of the partitions with the right symmetries under S_ℓ , these blocks should be considered indistinguishable. Therefore,

$$|\mathcal{C}_1| = \frac{1}{(\ell - 1)!} \tilde{A}_k^{(\ell-1)} (\dim \lambda)^2 , \quad (\text{A.10})$$

as there are $\dim \lambda$ Young tableaux of dimensionality $\dim \lambda$. For a fixed value of $\ell \geq 1$, the total number of trivial eigenvalues in $\tilde{\mathbf{T}}_L$ coming from all irreducible representations $\lambda \in S_\ell$ and divided by $\ell!$ is

$$\tilde{N}_{k1}(\ell) = \frac{1}{\ell!} \frac{1}{(\ell - 1)!} \tilde{A}_k^{(\ell-1)} \sum_{\lambda \in S_\ell} \dim^2 \lambda = \frac{1}{(\ell - 1)!} \tilde{A}_k^{(\ell-1)} , \quad (\text{A.11})$$

as $\sum_{\lambda \in S_\ell} \dim^2 \lambda = \ell!$. Finally, using (3.34) and after some algebra, it is not difficult to find the expression (A.2). ■

A.2 Extension of Lemma 4.2

Lemma A.2 *Fix $k \geq 1$, and $\ell = k$. Then, there are k non-trivial eigenvalues $\mu_{k,k,s}$ in the diagonal block $\tilde{\mathbf{T}}_{k,k}$ (once we have removed the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$). Each of these eigenvalues $\mu_{k,k,s}$ has multiplicity $k!$.*

PROOF. Let us first consider the following partition $(010'1')(22') \dots (kk')$ with $\ell = k$ links, where the primed numbers denote the sites on the top row, and the un-primed ones, those on the bottom row. This partition does not lead to the trivial eigenvalue $\mu_{k,k+1} = (-1)^k$, as it belongs to the class $\mathcal{C}_1^k$ defined in the proof of Lemma A.1 above. Then the action of the transfer matrix $\mathbf{T}_{k+1}$ [cf., (A.1)] on this partition generates exactly k partitions that can be

constructed as follows: starting from the basic partition $(011')(22') \dots (kk')$ with $\ell = k$ links, attach the site $0'$ to any of the k possible blocks. It is important to note that the ordering of the k links is preserved in the whole process. (Loosely speaking, there is “no room” to switch two links if we have $k + 1$ sites and k links.) This also means that the action of the transfer matrix $\mathbf{T}_{k+1}$ on any of these partitions is independent of the actual ordering of the links. Hence, for each of the $k!$ possible orderings of the k links, the transfer matrix will be the same (modulo some reordering of the partitions), and will have dimensionality k . Therefore, all irreducible representations of the symmetric group S_k will give the same k non-trivial eigenvalues, and the multiplicity of each of these eigenvalues in $\mathbf{T}_{k+1,k}$ is $\sum_{\lambda \in S_k} \dim^2 \lambda = k!$. ■

A.3 Extension of Corollary 4.3

Corollary A.3 *Fix $k \geq 1$. Then for all pairs (ℓ, λ) with $0 \leq \ell \leq k - 1$ and $\lambda \in S_\ell$, the diagonal block $\tilde{\mathbf{T}}_{k+1,\ell,\lambda}$ has $\tilde{N}_{k,0}(\ell)$ non-trivial eigenvalues $\mu_{k,\ell,\lambda,s}$ (once we have removed the trivial eigenvalue $\mu_{k+1,k} = (-1)^k$, which appears for every $1 \leq \ell \leq k + 1$). The value of $\tilde{N}_{k,0}(\ell)$ is given by:*

$$\tilde{N}_{k,0}(\ell) = \begin{cases} S_{k+1} + \sum_{p=0}^k \binom{k+1}{p+1} S_{k-p} & \text{if } \ell = 0 \\ Y_\ell \sum_{p=1}^k \binom{k}{p} \left\{ \begin{matrix} p \\ \ell \end{matrix} \right\} [S_{k-p} + S_{k+1-p}] & \text{if } 1 \leq \ell \leq k - 1 \\ k & \text{if } \ell = k \\ 0 & \text{if } \ell = k + 1 \end{cases} \quad (\text{A.12})$$

where S_n is given by (3.33), and Y_ℓ is the number of Young tableaux with ℓ cells.

PROOF. Let us first fix $0 \leq \ell < k$, and the irreducible representation λ of the symmetric group S_ℓ (of dimensionality $\dim \lambda$). The corresponding diagonal block is $\tilde{\mathbf{T}}_{k+1,\ell,\lambda}$, and its dimensionality is

$$\dim \tilde{\mathbf{T}}_{k+1,\ell,\lambda} = \frac{\tilde{N}_k(\ell)}{\ell!} \dim \lambda. \quad (\text{A.13})$$

The number of trivial eigenvalues $\mu_{k,k+1} = (-1)^k$ in $\tilde{\mathbf{T}}_{k+1,\ell,\lambda}$ is $\tilde{N}_{k,1}(\ell) \dim \lambda$ by Lemma A.1. Therefore, the number of non-trivial eigenvalues in this block is given by

$$\left[\frac{\tilde{N}_k(\ell)}{\ell!} - \tilde{N}_{k,1}(\ell) \right] \dim \lambda, \quad (\text{A.14})$$

and the number of non-trivial eigenvalues for all different values of $\lambda \in S_\ell$ is given by

$$\tilde{N}_{k,0}(\ell) = \left[\frac{\tilde{N}_k(\ell)}{\ell!} - \tilde{N}_{k,1}(\ell) \right] \sum_{\lambda \in S_\ell} \dim \lambda. \quad (\text{A.15})$$

This latter sum is equal to the number of Young tableaux with ℓ cells Y_ℓ (4.6).

We can provide a closed form for $\tilde{N}_{k,0}(\ell)$ for $0 \leq \ell < k$ by combining Lemma A.1, the definition $\tilde{N}_k(\ell) = |\tilde{A}_{k+1}^{(\ell)}|$, and (3.34)/(A.15). After some algebra we find Eq. (A.12) for $\ell < k$.

The case $\ell = k$ is derived directly from Lemma A.2. There are k non-trivial eigenvalues for every irreducible representation λ of S_k . Therefore, $\tilde{N}_{k,0}(k) = k$.

Finally, for $\ell = k + 1$, the number of non-trivial eigenvalues is zero, as all eigenvalues are trivial in this sector. Note that $\tilde{N}_k(k + 1) = (k + 1)!$ [cf. (3.34)], and that $\tilde{N}_{k,1}(k + 1) = 1$ [cf. (4.4)], so that $\tilde{N}_{k,0}(k + 1) = 0$ by Eq. (A.15). ■

A.4 Proof of Lemma 4.6

All the matrix elements of $\tilde{T}_{k+1,\ell,\lambda}$ are polynomials in Q . Therefore, $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ will be a polynomial in Q . However, not all the matrix elements of $\hat{T}_{k+1,\ell,\lambda}$ are polynomials in Q ; so we cannot conclude that $\text{tr } \hat{T}_{k+1,\ell,\lambda}^n$ is a polynomial in Q .

We need a different argument. Let us consider any partition $\mathcal{P}$ belonging to $\tilde{\mathcal{A}}_{k+1}^{(\ell)}$. Then, its contribution to $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ will be the sum of the contributions of all the diagrams that start with the partition $\mathcal{P}$ and end with the same partition after n steps. By inspection of the form of the transfer matrix T_{k+1} (3.8) and its components V_i and H_{0i} (3.7), it is clear that there is a contribution proportional to $(-Q)^{2kn}$ due to the application of all the identity operators in T_{k+1} (for each layer, there are $3k$ of them, and only $2k$ take the factor $-Q$). Indeed, this is the minimum number of $-Q$ factors one can possibly obtain for a diagram of this type. Therefore, the minimum power of Q that appears in $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ is Q^{2kn} , which is exactly the prefactor in the definition of $\hat{T}_{k+1}$ (3.19)/(3.20). In conclusion, $\text{tr } \hat{T}_{k+1,\ell,\lambda}^n$ is indeed a polynomial in Q .

Once the polynomial character of $\text{tr } \hat{T}_{k+1,\ell,\lambda}^n$ is established, we have to take care of the degree of this polynomial. The lemma is proved if we are able to show that all the entries in $\tilde{T}_{k+1,\ell,\lambda}$ have powers of Q of degree at most $3k + \min(1 - \ell, 0)$. This would imply, that $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ is a polynomial in Q of degree at most $n[3k + \min(1 - \ell, 0)]$, and that $\text{tr } \hat{T}_{k+1,\ell,\lambda}^n$ is a polynomial in Q of degree at most $n[k + \min(1 - \ell, 0)]$, as claimed.

Let us first consider the case $\ell = 0$. Let us start with an arbitrary partition of the top row $\mathcal{P} \in \tilde{\mathcal{A}}_{k+1}^{(0)}$. When we apply $\tilde{T}_{k+1}$ to that partition, the maximum number of $-Q$ factors that can appear is $(-Q)^{3k}$: we apply the part $-Q\mathbf{1}$ for each vertical operator V_i , and the join operator $-QJ_{0i}$ for each horizontal operator H_{0i} . We obtain for all $\mathcal{P} \in \tilde{\mathcal{A}}_{k+1}^{(0)}$ the same final partition: $\{\{0, 1, \dots, k\}\}$. Therefore, $\text{tr } \tilde{T}_{k+1,0,\lambda}^n$ is a polynomial in Q of degree at most $3kn$.

The case $\ell = 1$ is similar. If we start with an arbitrary partition of the top row $\mathcal{P} \in \tilde{\mathcal{A}}_{k+1}^{(1)}$ with 0 marked, and we apply the same operators as above, we end up with the partition $\{\{\overline{0}, \overline{1}, \dots, \overline{k}\}\}$ (where the overline over a site denotes that this site is connected to the bottom row). The coefficient corresponding to this partition is a polynomial in Q of degree at most $3k$. Therefore, $\text{tr } \tilde{T}_{k+1,1,\lambda}^n$ is a polynomial in Q of degree at most $3kn$.

The case $2 \leq \ell \leq k + 1$ is similar. Let us start with the following simple partition in $\tilde{\mathcal{A}}_{k+1}^{(\ell)}$: $\{\{\overline{0}, \overline{\ell}, \dots, \overline{k}\}, \{\overline{1}\}, \dots, \{\overline{\ell - 1}\}\}$. The argument is similar to the previous cases, except that

we cannot join the site 0 to any of the other $\ell - 1$ blocks; since if we did, then the number of links would be smaller than ℓ . Therefore, we get a diagonal entry which is a polynomial in Q of degree at most $3k - (\ell - 1)$. Therefore, $\text{tr } \tilde{T}_{k+1,\ell,\lambda}^n$ is a polynomial in Q of degree at most $n[3k - (\ell - 1)]$. ■

Remark. Note that the above Lemma for $k = \ell + 1$ implies that the trace $\text{tr } \tilde{T}_{k+1,k+1,\lambda}^n$ is a polynomial in Q of degree at most $n[3k - k] = 2nk$. Thus, $\text{tr } \hat{T}_{k+1,k+1,\lambda}^n$ is a polynomial in Q of degree at most $2nk - 2nk = 0$, in agreement with Lemma A.1.

B The polynomial $\Phi_{G(119,7)}$

i	a_i
0	240453758183717079931230416441214627161100181583221695778758847017660
1	3778010581676303383947166862404894626185168386864045862610150115812052
2	30335569899732630785396756910315613411372095140384361997526113182374163
3	165825101972051263346220423069919998872239305113596582973635632055394977
4	693579540994844285783536577772823633624035569377766733899721814879025261
5	2365298420910039361031441041999881902182643546761840393901756834507719194
6	6843881667041711337123509683076504016112680686224789087846006938716709936
7	17263457862032800683223779137831846913199242114984324936441639648268053289
8	38713433713214705146001095156944656719927970414458008688062316276391488388
9	78323139898934533680532026518046626525971702819240021820160355852376101900
10	144596924706446634397013884950643553134413962554815223156161387400333508225
11	245805490880525223783583677086377232647690995050227537846281841251646914824
12	387578029394911704914155641300482722338098395075782035706404804765273096365
13	570251442350241862197997781738305884386349966454124613787648726411773264858
14	786830079084979854704322218255681568053263515605219068314256695784834361687
15	1022415665287852311682199189937274383849816815595817077282491287814210089327
16	1255617405561053292049115611630960519206807161888631329237623843787650902598
17	1461844034312439449575494091690068905133069052914279852509149780627087441657
18	1617740821249798765950531657277432254427341252461560806920840933023592630423
19	1705621288663026987807981694789591864355698512752538498113371227662898599836
20	1716721077841948079359918580031477807370750494656542267873960267539031216693
21	1652471749735377739204603329938965183162646296909313884512941374410264086521
22	1523600861689858982867164652432211354402872413669859092087649541160917763276
23	1347478648579683679991904590350043152972817068987716895096810710111624258414
24	1144543577841773436389066666439373326178625474359360165078175860151437888418
25	934744903425435172192816740334988586706047594412696933301815485640045053088
26	734761463017109267283189805699271455927373168195072916098687225728186938239
27	556406139434285183296690956661070857540502747488343675526234531927677714022

Table 2: Coefficients of the flow polynomial $\Phi_{G(119,7)}(Q) = (Q - 1)(Q - 2)(Q - 3) \sum_{i=0}^{117} (-1)^{i+1} a_i Q^i$.

i	a_i
28	406248919977322871211475078592598791327840251152453917692944363153498710100
29	286204685555683178005609409552066982429730669340777755008844846916076522994
30	194690835588052864277724799113473665492499996408415866330514336755095511519
31	127959409173717651616971568749651845945079328780268936837379415435944126802
32	81303058361075417592304554906219204545070829443708169095378279034181616797
33	49966395799032436266189002396534646533235474102922584729978792549768709570
34	29716297381875173070101426039892101696811371497930415707024934873184994971
35	17109926011423464771467167427507814898291755276949983853151230858292134005
36	9541424411645170221820026129682497673840857753028326218052776323947917867
37	5155250259203698010398273841766642027281260863127250587163264787844562005
38	2699625968148386503782707617994694138490854643501950268752229332983496484
39	1370590172276847791788823798202187745155060493411059843973697377813429115
40	674811604481186691941181395279228274700479647745836519066846509110686871
41	322283263195622661124593258235203925484545860053525952388302407768566612
42	149338540704756141327132733140880585247255726988170348306522536601036428
43	67154254327764896250779574488797365684816293909986077788680077649219059
44	29310387712861712336349988437753297027267572645934195793910184667409525
45	12418975519744799441679142496250486367091770751393334192303593690937216
46	5108897246470415157235242805025854028977310258147102091244333174463070
47	2040790191809533248688898356534425242558953177861065156649563146370679
48	791669367678994305215119572575588439395174310193863932545018547493949
49	298263611514781343272224652335166519468558609433860386772388696144855
50	109143202315633324987330964029233262735802381917881292972386146290583
51	38793120529207946517836936596295050010421179961661603601012761033976
52	13393333822523527127943242411311403493467668611668190366289160119798
53	4491649304261905729281333948467503543032844169847443690259485449787
54	1463208911995358578670442816667443146618570037914577188051167519252
55	463002576567921398643877003491825254798116352410207026582354227639
56	142306049793794840801481088729855032253716088115496415256458286993
57	42482066138212698853729014913405235508231962094892775535623301954
58	12316937820630034457717661311813898450293797908401593794293724758
59	3468010035454592592294532401090077198723295654493559716526595085
60	948193746090263874793189178585310649056926723273402296835255341
61	251712395999861219430022121008687006528308619159639108633408792
62	64870522436154805702181162937007326080100049608985554185883024
63	16227914273399692643538894326321813038077624088578343763024693
64	3939856175099340812778271579647788724054817562072496203271622
65	928159554704604760538186427442530041159446537967362956747430
66	212130925228863323822024721431112601866613699451131860235710
67	47025221622368034908085340305586428332317539464533841648123
68	10108841512034189983877166351043431883332647922883010856327
69	2106711947665769238334508701464763203793865636828903442294
70	425522761043753110078251752857561183613194192306420512412
71	83277105914548332097899604836384635641176843712157856926
72	15786114232928385188569755800951974220820338221822797673
73	2897497349085917262770771233534175446735608061805601330

Table 2: (Continued.)

i	a_i
74	514765848404059046836837761417162573532059958487097353
75	88483984104925516580422896030293322509203586666446608
76	14709751385762293879136933265239581784434168823726105
77	2363940443321701747333111550458096052434867769928216
78	367070636910713149669510355126240632283426228583990
79	55045491030169621973045797552387375721405805002210
80	7967363593383576742789482483060195429602666353792
81	1112432655029526956212696750602542449173138786358
82	149736221917777447936668185649515326795606657139
83	19417037687233804941366229781880098811952696949
84	2423988165427613282917222688678473423362471885
85	291095664115059080227254369870423630080267269
86	33600127040413356951575245326064861734434090
87	3724448845910258224439985334986604271814630
88	396083067053161605214005007534380917557302
89	40370994515077502586895936288134103918883
90	3939422436008602219258246065351204112198
91	367585380265857178359590490307505526698
92	32755670407906919539877604731444999292
93	2783619278733903044429223967432069173
94	225251413853620678004411181186579261
95	17327620037637857872926704478452708
96	1264839938089389154908070831758552
97	87436328017906227077510426235281
98	5711550730808768130994711119064
99	351695047292333601570905687819
100	20358915651379087237820736779
101	1104604810216428455300363966
102	55981869638360027493471673
103	2640008664516116400095478
104	115338644735668374521021
105	4644751096461226418781
106	171402937526338140238
107	5756292666737489361
108	174485239351077202
109	4726362454421551
110	113000004047458
111	2347463061581
112	41510490693
113	607499109
114	7064107
115	61191
116	351
117	1

Table 2: (Continued.)

Acknowledgments

We wish to thank Gordon Royle for suggesting us to study the generalised Petersen graphs, and Alan Sokal for helpful conversations and for a critical reading of the manuscript that gave rise to many constructive comments.

Both authors also thank the Isaac Newton Institute for Mathematical Sciences, University of Cambridge, for hospitality during the programme on Combinatorics and Statistical Mechanics (January–June 2008), where this project started. JS also thanks the École Normale Supérieure for hospitality in June 2009.

The research of JS was supported in part by Spanish MEC grants FPA2009-08785 and MTM2008-03020, and by U.S. National Science Foundation grant PHY-0424082. The research of JLJ was supported in part by the European Community Network ENRAGE (grant MRTN-CT-2004-005616), and by the Agence Nationale de la Recherche (grant ANR-06-BLAN-0124-03)

References

- [1] K.I. Appel and W. Haken, *Every planar map is four colorable*, Illinois J. Math. **21**, 429–567 (1977).
- [2] R.J. Baxter, *Potts model at the critical temperature*, J. Phys. C: Solid State Phys. **6**, L445–L448 (1973).
- [3] R.J. Baxter, H.N.V. Temperley and S.E. Ashley, *Triangular Potts model at its transition temperature, and related models*, Proc. Roy. Soc. London A **358**, 535–559 (1978).
- [4] R.J. Baxter, *Critical antiferromagnetic square-lattice Potts model*, Proc. Roy. Soc. London A **383**, 43–54 (1982).
- [5] R.J. Baxter, *q colourings of the triangular lattice*, J. Phys. A: Math. Gen. **19**, 2821–2839 (1986).
- [6] R.J. Baxter, *Chromatic polynomials of large triangular lattices*, J. Phys. A: Math. Gen. **20**, 5241–5261 (1987).
- [7] A. Bedini and J.L. Jacobsen, *A tree-decomposed transfer matrix for computing exact Potts model partition functions for arbitrary graphs, with applications to planar graph colourings*, J. Phys. A: Math. Theor. **43**, 385001 (2010), arXiv:1003.4847.
- [8] S. Beraha and J. Kahane, *Is the four-color conjecture almost false?*, J. Combin. Theory Ser. B **27**, 1–12 (1979).
- [9] S. Beraha, J. Kahane and N.J. Weiss, *Limits of zeroes of recursively defined polynomials*, Proc. Nat. Acad. Sci. USA **72**, 4209 (1975).
- [10] S. Beraha, J. Kahane and N.J. Weiss, *Limits of zeroes of recursively defined families of polynomials*, in *Studies in Foundations and Combinatorics* (Advances in Mathematics Supplementary Studies, Vol. 1), ed. G.-C. Rota, pp. 213–232 (Academic Press, New York, 1978).

- [11] S. Beraha, J. Kahane and N.J. Weiss, *Limits of chromatic zeros of some families of maps*, J. Combin. Theory Ser. B **28**, 52–65 (1980).
- [12] D.A. Bini and G. Fiorentino, Numerical computation of polynomial roots using MPSolve version 2.2 (January 2000). Software package and documentation available for download at <ftp://ftp.dm.unipi.it/pub/mpsolve/MPSolve-2.2.tgz>
- [13] D.A. Bini and G. Fiorentino, *Design, analysis, and implementation of a multiprecision polynomial rootfinder*, Num. Algorithms **23**, 127–173 (2000).
- [14] G.D. Birkhoff, *A determinant formula for the number of ways of coloring a map*, Ann. Math. **14**, 42–46 (1912).
- [15] G.D. Birkhoff and D.C. Lewis, *Chromatic polynomials*, Trans. Amer. Math. Soc. **60**, 355–451 (1946).
- [16] F. Castagna and G. Prins, *Every generalized Petersen graph has a Tait coloring*, Pacific J. Math. **40**, 53–58 (1972).
- [17] R. Diestel, *Graph Theory*, (Springer Verlag, Heidelberg, New York, 2005).
- [18] S.-C. Chang, J. Salas and R. Shrock, *Exact Potts model partition functions for strips of the square lattice*, J. Stat. Phys. **107**, 1207–1253 (2002), arXiv:cond-mat/0108144.
- [19] Ph. Flajolet and R. Sedgewick, *Analytic Combinatorics*, (Cambridge University Press, Cambridge, 2009).
- [20] C.M. Fortuin and P.W. Kasteleyn, *On the random-cluster model: I. Introduction and relation to other models*, Physica **57**, 536–564 (1972).
- [21] R.L. Graham, D.E. Knuth, and O. Patashnik, *Concrete Mathematics*, 2nd edition (Addison–Wesley, 1994).
- [22] G. Haggard, D.J. Pearce, and G.F. Royle, *Computing Tutte polynomials*, ACM Transactions on Mathematical Software, (to appear).
- [23] T. Halverson and A. Ram, *Partition algebras*, Eur. J. Comb. **26**, 869–921 (2005).
- [24] B. Jackson, *A zero-free interval for flow polynomials of cubic graphs*, J. Comb. Theory B **97**, 127–143 (2007).
- [25] B. Jackson, *Zeros of chromatic and flow polynomials of graphs*, J. Geom. **76**, 95–109 (2003), arXiv:math/0205047.
- [26] J.L. Jacobsen and J. Salas, *Transfer matrices and partition-function zeros for antiferromagnetic Potts models. IV. Chromatic polynomial with cyclic boundary conditions*, J. Stat. Phys. **122**, 705–760 (2006), arXiv:cond-mat/0407444.
- [27] J.L. Jacobsen and J. Salas, *Phase diagram of the chromatic polynomial on a torus*, Nucl. Phys. B **783**, 238–296 (2007), arXiv:cond-mat/0703228.

- [28] J.L. Jacobsen, J. Salas and A.D. Sokal, *Transfer matrices and partition-function zeros for antiferromagnetic Potts models. III. Triangular-lattice chromatic polynomial*, J. Stat. Phys. **112**, 921–1017 (2003), arXiv:cond-mat/0204587.
- [29] J.L. Jacobsen and H. Saleur, *The antiferromagnetic transition for the square-lattice Potts model*, Nucl. Phys. B **743**, 207–248 (2006), arXiv:cond-mat/0512058.
- [30] J.L. Jacobsen and H. Saleur, *Boundary chromatic polynomial*, J. Stat. Phys. **132**, 707–719 (2008), arXiv:0803.2665.
- [31] J.L. Jacobsen and H. Saleur, *Combinatorial aspects of boundary loop models*, J. Stat. Mech. P01021 (2008).
- [32] F. Jaeger, *Flows and generalized coloring theorems in graphs*, J. Comb. Theory B **26**, 205–216 (1979).
- [33] F. Jaeger, *Nowhere-zero flow problems*, in L.W. Beineke and R.J. Wilson (eds.), *Selected topics in graph theory 3* (Academic Press, 1988).
- [34] G. Kirchhoff, *Ueber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird*, Annalen der Physik und Chemie **148**, 497–508 (1847).
- [35] M. Kochol, *Restrictions on the smallest counterexamples to the 5-flow conjecture*, Combinatorica **26**, 83–89 (2006).
- [36] P.P. Martin, *The structure of the partition algebras*, J. Algebra **183**, 319–358 (1996).
- [37] J.G. Oxley, *Colourings, packing and the critical problem*, Quart. J. Math. Oxford **29**, 11–22 (1978).
- [38] V. Pasquier and H. Saleur, *Common structures between finite systems and conformal field theories through quantum groups*, Nucl. Phys. B **330**, 523–556 (1990).
- [39] R.B. Potts, *Some generalized order-disorder transformations*, Proc. Camb. Phil. Soc. **48**, 106–109 (1952).
- [40] J.-F. Richard and J.L. Jacobsen, *Eigenvalue amplitudes of the Potts model on a torus*, Nucl. Phys. B **769**, 256–274 (2007), arXiv:math-ph/0608055.
- [41] G.F. Royle, *Planar triangulations with real chromatic roots arbitrarily close to four*, Annals Combin. **12**, 195–210 (2008), arXiv:math/0511304.
- [42] B.E. Sagan, *The symmetric group: Representations, combinatorial algorithms, and symmetric functions*, Graduate Texts in Mathematics **203** (Springer-Verlag, 2000).
- [43] J. Salas and A.D. Sokal, *Transfer matrices and partition-function zeros for antiferromagnetic Potts models. VI. Square lattice with special boundary conditions*, arXiv:1002.3761.
- [44] H. Saleur, *Zeroes of chromatic polynomials: a new approach to Beraha conjecture using quantum groups*, Comm. Math. Phys. **132**, 657–679 (1990).

- [45] H. Saleur, *The antiferromagnetic Potts model in two dimensions: Berker-Kadanoff phase, antiferromagnetic transition, and the role of Beraha numbers*, Nucl. Phys. B **360**, 219–263 (1991).
- [46] P.D. Seymour, *Nowhere-zero 6-flows*, J. Comb. Theory B **30**, 130–135 (1981).
- [47] N.J.A. Sloane, Sloane’s On-Line Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences/index.html>.
- [48] A.D. Sokal, *Chromatic roots are dense in the whole complex plane*, Comb. Prob. Comp. **13**, 221–261 (2004), arXiv:cond-mat/0012369.
- [49] R. Steinberg, *Tutte’s 5-flow conjecture for the projective plane*, J. Graph Theory **8**, 277–289 (1984).
- [50] C. Thomassen, *The zero-free intervals for chromatic polynomials of graphs*, Combin. Probab. Comput. **6**, 497–442 (1997).
- [51] W.T. Tutte, *A ring in graph theory*, Proc. Camb. Phil. Soc. **43**, 26–40 (1947).
- [52] W.T. Tutte, *On the imbedding of linear graphs in surfaces*, Proc. London Math. Soc. **51**, 474–483 (1950).
- [53] W.T. Tutte, *A contribution to the theory of chromatic polynomials*, Can. J. Math. **6**, 81–90 (1954).
- [54] W.T. Tutte, *Graph theory* (Addison-Wesley, New York, 1984).
- [55] C.D. Wakelin, *Chromatic Polynomials*, Ph. D. thesis, University of Nottingham, 1994.
- [56] M.E. Watkins, *A theorem on Tait colorings with an application to the generalized Petersen graphs*, J. Combin. Theory **6**, 152–164 (1969).
- [57] D.J.A. Welsh, <http://garden.irmacs.sfu.ca/?q=category/welsh>. Also cited in Ref. [22, Conjecture 1] and in Ref. [25, Conjecture 33].
- [58] H. Whitney, *The coloring of graphs*, Ann. Math. **33**, 688–718 (1932).
- [59] H. Whitney, *A logical expansion in mathematics*, Bull. Amer. Math. Soc. **38**, 572–580 (1932).
- [60] D.R. Woodall, *The largest real zero of the chromatic polynomial*, Discrete Math. **101**, 333–341 (1997).
- [61] F.Y. Wu, *Potts model and graph theory*, J. Stat. Phys. **52**, 99–112 (1988).
- [62] F.Y. Wu and Y.K. Wang, *Duality transformation in a many-component spin model*, J. Math. Phys. **17**, 439 (1976).
- [63] C.-Q. Zhang, *Integer flows and cycle covers of graphs* (Marcel Dekker, New York, 1997).